



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁷ : C07D 401/14, A61K 31/40, C07D 401/12, 413/14</p>	<p>A1</p>	<p>(11) International Publication Number: WO 00/04013</p> <p>(43) International Publication Date: 27 January 2000 (27.01.00)</p>
<p>(21) International Application Number: PCT/US99/15581</p> <p>(22) International Filing Date: 9 July 1999 (09.07.99)</p> <p>(30) Priority Data: 09/115.497 14 July 1998 (14.07.98) US</p> <p>(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application US 09/115,497 (CON) Filed on 14 July 1998 (14.07.98)</p> <p>(71) Applicant (for all designated States except US): MERCK & CO., INC. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(72) Inventors; and</p> <p>(75) Inventors/Applicants (for US only): GOULET, Mark [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). ASHTON, Wallace, T. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). CHU, Lin [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). FISHER, Michael, H. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). LIN, Peter [GB/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). PONPIPOM, Mitree, M.</p>		<p>[US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). WYVRATT, Matthew, J. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). GIROTRA, Narindar, N. [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US). YOUNG, Jonathan [US/US]; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(74) Common Representative: MERCK & CO., INC.; 126 East Lincoln Avenue, Rahway, NJ 07065 (US).</p> <p>(81) Designated States: AE, AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GD, GE, HR, HU, ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ZA, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>
<p>(54) Title: ANTAGONISTS OF GONADOTROPIN RELEASING HORMONE</p> <div style="text-align: center; margin: 20px 0;"> <p style="text-align: right;">(I)</p> </div>		
<p>(57) Abstract</p> <p>The present invention relates to compounds of formula (I) which are non-peptide antagonists of GnRH which can be used to treat a variety of sex-hormone related conditions in men and women, to methods for their preparation, and to methods and pharmaceutical compositions containing said compounds for use in mammals.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

TITLE OF THE INVENTION

ANTAGONISTS OF GONADOTROPIN RELEASING HORMONE

5 This application is a continuation-in-part of copending application Serial No. 08/760,186, filed December 5, 1996.

BACKGROUND OF THE INVENTION

10 The gonadotropin-releasing hormone (GnRH), also referred to as luteinizing hormone-releasing hormone (LHRH), is a decapeptide that plays a key role in human reproduction. The hormone is released from the hypothalamus and acts on the pituitary gland to stimulate the biosynthesis and secretion of luteinizing hormone (LH) and follicle-stimulating hormone (FSH). LH released from the pituitary gland is primarily responsible for the regulation of gonadal steroid production in both sexes, whereas FSH regulates spermatogenesis in males and follicular development in females. GnRH agonists and antagonists have proven effective in the treatment of certain conditions which require inhibition of LH/FSH release. In particular, GnRH-based therapies have proven effective in the treatment of endometriosis, uterine fibroids, polycystic ovarian disease, precocious puberty and several gonadal steroid-dependent neoplasia, most notably cancers of the prostate, breast and ovary. GnRH agonists and antagonists have also been utilized in various assisted fertilization techniques and have been investigated as a potential contraceptive in both men and women. They have also shown possible utility in the treatment of pituitary gonadotrophe adenomas, sleep disorders such as sleep apnea, irritable bowel syndrome, premenstrual syndrome, benign prostatic hyperplasia, hirsutism, as an adjunct to growth hormone therapy in growth hormone deficient children, and in murine models of lupus.

30 Current GnRH antagonists are GnRH-like decapeptides which are generally administered intravenously or subcutaneously presumably because of negligible oral activity. These have amino acid substitutions usually at positions one, two, three, six and ten.

35 Non-peptide GnRH antagonists offer the possible advantage of oral administration. Non-peptide GnRH antagonists have been

described in European Application 0 219 292 and in De, B. et al., J. Med. Chem., **32**, 2036-2038 (1989), in WO 95/28405, WO 95/29900 and EP 0679642 all to Takeda Chemical Industries, Ltd.

5 Substituted indoles known in the art include those described in the following patents and patent applications. US Patent No. 5,030,640 discloses alpha-heterocyclic ethanol aminoalkyl indoles which are potent β -agonists. US Patent No. 4,544,663 discloses indolamine derivatives which are allegedly useful as male anti-fertility agents. WO 90/05721 discloses alpha-amino-indole-3-acetic acids useful as
10 anti-diabetic, anti-obesity and anti-atherosclerotic agents. French patent 2,181,559 discloses indole derivatives with sedative, neuroleptic, analgesic, hypotensive, antiserotonin and adrenolytic activity. Belgian patent 879381 discloses 3-aminoalkyl-1H-indole-5-thioamide and carboxamide derivatives as cardiovascular agents used to treat
15 hypertension, Raynaud's disease and migraine. WO 97/21435, WO 97/21703, WO 97/21707 and WO 97/21704 disclose non-peptidyl, indole derivatives as GnRH antagonists.

SUMMARY OF THE INVENTION

20 The present invention relates to compounds which are non-peptide antagonists of GnRH which can be used to treat a variety of sex-hormone related conditions in men and women, to methods for their preparation, and to methods and pharmaceutical compositions containing said compounds for use in mammals.

25 Because of their activity as antagonists of the hormone GnRH, the compounds of the present invention are useful to treat a variety of sex-hormone related conditions in both men and women. These conditions include endometriosis, uterine fibroids, polycystic ovarian disease, hirsutism, precocious puberty, gonadal steroid-
30 dependent neoplasias such as cancers of the prostate, breast and ovary, gonadotrophe pituitary adenomas, sleep apnea, irritable bowel syndrome, premenstrual syndrome and benign prostatic hypertrophy. They are also useful as an adjunct to treatment of growth hormone deficiency and short stature, and for the treatment of systemic lupus
35 erythematosus. Further, the compounds of the invention may be useful

in *in vitro* fertilization and as contraceptives. The compounds may also be useful in combination with androgens, estrogens, progesterones, antiestrogens and antiprogestogens for the treatment of endometriosis, fibroids and in contraception. They may also be useful in combination with testosterone or other androgens or antiprogestogens in men as a contraceptive. The compounds may also be used in combination with an angiotensin-converting enzyme inhibitor such as Enalapril or Captopril, an angiotensin II-receptor antagonist such as Losartan or a renin inhibitor for the treatment of uterine fibroids. Additionally, the compounds of the invention may also be used in combination with bisphosphonates (bisphosphonic acids) and other agents, such as growth hormone secretagogues, e.g. MK-0677, for the treatment and prevention of disturbances of calcium, phosphate and bone metabolism, in particular, for the prevention of bone loss during therapy with the GnRH antagonist, and in combination with estrogens, progesterones, antiestrogens, antiprogestins and/or androgens for the prevention or treatment of bone loss or hypogonadal symptoms such as hot flashes during therapy with the GnRH antagonist.

Additionally, a compound of the present invention may be co-administered with a 5 α -reductase 2 inhibitor, such as finasteride or epristeride; a 5 α -reductase 1 inhibitor such as 4,7b-dimethyl-4-aza-5 α -cholestan-3-one, 3-oxo-4-aza-4,7b-dimethyl-16b-(4-chlorophenoxy)-5 α -androstane, and 3-oxo-4-aza-4,7b-dimethyl-16b-(phenoxy)-5 α -androstane as disclosed in WO 93/23420 and WO 95/11254; dual inhibitors of 5 α -reductase 1 and 5 α -reductase 2 such as 3-oxo-4-aza-17b-(2,5-trifluoromethylphenyl-carbamoyl)-5 α -androstane as disclosed in WO 95/07927; antiandrogens such as flutamide, casodex and cyproterone acetate, and alpha-1 blockers such as prazosin, terazosin, doxazosin, tamsulosin, and alfuzosin.


Further, a compound of the present invention may be used in combination with growth hormone, growth hormone releasing hormone or growth hormone secretagogues, to delay puberty in growth hormone deficient children, which will allow them to continue to gain height before fusion of the epiphyses and cessation of growth at puberty.

Further, a compound of the present invention may be

used in combination or co-administered with a compound having luteinizing hormone releasing activity such as a peptide or natural hormone or analog thereof. Such peptide compounds include leuporelin, gonadorelin, buserelin, triptorelin, goserelin, nafarelin, histrelin, deslorelin, meterlin and recirelin.

DETAILED DESCRIPTION OF THE INVENTION


The present invention relates to compounds of the general formula



wherein

A is

C1-C6 alkyl, substituted C1-C6 alkyl, C3-C7 cycloalkyl, substituted C3-C7 cycloalkyl, C3-C6 alkenyl, substituted C3-C6 alkenyl, C3-C6 alkynyl, substituted C3-C6 alkynyl, C1-C6 alkoxy, or C0-C5 alkyl-S(O)_n-C0-C5 alkyl, C0-C5 alkyl-O-C0-C5 alkyl, C0-C5 alkyl-NR₁₈-C0-C5 alkyl where R₁₈ and the




C0-C5 alkyl can be joined to form a ring, or a single bond;


R₀ is

hydrogen, C1-C6 alkyl, substituted C1-C6 alkyl, wherein the substituents are as defined below; aryl, substituted aryl, aralkyl or substituted aralkyl, wherein the substituents are as defined for R₃, R₄ and R₅;

R₁ is



- the nitrogen atoms contained in the R₁ heteroaromatic rings may exist
 either as drawn or, when chemically allowed, in their
 oxidized (N→O, N-OH) state;
- 5 R₂ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aralkyl,
 substituted aralkyl, aryl, substituted aryl, alkyl -OR₁₁,
 C₁-C₆(NR₁₁R₁₂), C₁-C₆(CONR₁₁R₁₂) or C(NR₁₁R₁₂)NH;
- R₂ and A taken together form a ring of 5-7 atoms;
- 10 R₃, R₄ and R₅ are independently hydrogen, C₁-C₆ alkyl, substituted
 C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, CN,

- nitro, C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, aryl, substituted aryl, aralkyl, substituted aralkyl, R₁₁O(CH₂)_p-, R₁₁C(O)O(CH₂)_p-, R₁₁OC(O)(CH₂)_p-, -(CH₂)_pS(O)_nR₁₇-, (CH₂)_pC(O)NR₁₁R₁₂ or halogen; wherein R₁₇ is hydrogen, C₁-C₆ alkyl, C₁-C₃ perfluoroalkyl, aryl or substituted aryl;
- 5 R₃ and R₄ taken together form a carbocyclic ring of 3-7 carbon atoms or a heterocyclic ring containing 1-3 heteroatoms selected from N, O and S;
- 10 R₆ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, C₁-C₃ perfluoroalkyl, CN, NO₂, halogen, R₁₁O(CH₂)_p-, NR₂₁C(O)R₂₀, NR₂₁C(O)NR₂₀R₂₁ or SO_nR₂₀;
- R₇ is hydrogen, C₁-C₆ alkyl, or substituted C₁-C₆ alkyl, unless X is hydrogen or halogen, then R₇ is absent;
- 15 R₈ is C(O)OR₂₀, C(O)NR₂₀R₂₁, NR₂₀R₂₁, C(O)R₂₀, NR₂₁C(O)R₂₀, NR₂₁C(O)NR₂₀R₂₁, NR₂₀S(O)₂R₂₁, NR₂₁S(O)₂NR₂₀R₂₁, OC(O)R₂₀, OC(O)NR₂₀R₂₁, OR₂₀, SO_nR₂₀, S(O)_nNR₂₀R₂₁, a heterocyclic ring or bicyclic heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅, C₁-C₆ alkyl or substituted C₁-C₆ alkyl; or
- 20 R₇ and R₈ taken together form a heterocyclic ring containing one or more heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅;
- 25 R₉ and R_{9a} are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl; aryl or substituted aryl, aralkyl or substituted aralkyl when m≠0; or
- R₉ and R_{9a} taken together form a carbocyclic ring of 3-7 atoms or  when m≠0;
- 30 R₉ and A taken together form a heterocyclic ring containing 3-7 carbon atoms and one or more heteroatoms when m≠0; or
- R₁₀ and R_{10a} are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl; or

- R₁₀ and R_{10a} taken together form a carbocyclic ring of 3-7 atoms or $\text{O}=\text{C}$;
 R₉ and R₁₀ taken together form a carbocyclic ring of 3-7 carbon atoms or
 a heterocyclic ring containing one or more heteroatoms
 when m≠0; or
- 5 R₉ and R₂ taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms when m≠0; or
 R₁₀ and R₂ taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms;
- 10 R₁₀ and A taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms; or
 R₁₁ and R₁₂ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-
 C₆ alkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, a
 carbocyclic ring of 3-7 atoms or a substituted carbocyclic ring
 containing 3-7 atoms;
- 15 R₁₁ and R₁₂ taken together can form an optionally substituted ring of 3-7
 atoms;
- R₁₃ is hydrogen, OH, NR₇R₈, NR₁₁SO₂(C₁-C₆ alkyl),
 NR₁₁SO₂(substituted C₁-C₆ alkyl), NR₁₁SO₂(aryl),
 NR₁₁SO₂(substituted aryl), NR₁₁SO₂(C₁-C₃ perfluoroalkyl);
 20 SO₂NR₁₁(C₁-C₆ alkyl), SO₂NR₁₁(substituted C₁-C₆ alkyl),
 SO₂NR₁₁(aryl), SO₂NR₁₁(substituted aryl), SO₂NR₁₁(C₁-C₃
 perfluoroalkyl); SO₂NR₁₁(C(O)C₁-C₆ alkyl); SO₂NR₁₁(C(O)-
 substituted C₁-C₆ alkyl); SO₂NR₁₁(C(O)-aryl);
 SO₂NR₁₁(C(O)-substituted aryl); S(O)_n(C₁-C₆ alkyl); S(O)_n
 25 (substituted C₁-C₆ alkyl), S(O)_n(aryl), S(O)_n(substituted
 aryl), C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, C₁-C₆
 alkoxy, substituted C₁-C₆ alkoxy, COOH, halogen, NO₂ or
 CN;
- R₁₄ and R₁₅ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆
 30 alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, CN, nitro,
 C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, aryl,
 substituted aryl, aralkyl, substituted aralkyl, R₁₁O(CH₂)_p-,
 R₁₁C(O)O(CH₂)_p-, R₁₁OC(O)(CH₂)_p-, -(CH₂)_pS(O)_nR₁₇, -

- (CH₂)_pC(O)NR₁₁R₁₂ or halogen; wherein R₁₇ is hydrogen, C₁-C₆ alkyl, C₁-C₃ perfluoroalkyl, aryl or substituted aryl;
- R₁₆ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, or N(R₁₁R₁₂);
- 5 R₁₈ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C(O)OR₁₁, C(O)NR₁₁R₁₂, C(O)R₁₁, S(O)_nR₁₁;
- R₁₉ is either the definition of R₁₃ or R₁₄;
- R₂₀ and R₂₁ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, a
- 10 carbocyclic ring of 3-7 atoms, a substituted carbocyclic ring containing 3-7 atoms, a heterocyclic ring or bicyclic heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅, C₁-C₆-alkyl substituted by a heterocyclic ring or bicyclic
- 15 heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅;
- R₂₀ and R₂₁ taken together can form an optionally substituted ring of 3-7 atoms;
- 20 X is N, O, S(O)_n, C(O), (CR₁₁R₁₂)_p, a single bond to R₈, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, or substituted C₂-C₆ alkynyl; when X is O, S(O)_n, C(O), or CR₁₁R₁₂ only R₈ is possible;
- Z is O, S or NR₁₁;
- 25 m is 0-3;
- n is 0-2;
- p is 0-4; and
- 30 the alkyl, cycloalkyl, alkenyl and alkynyl substituents are selected from C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, hydroxy, oxo, cyano, C₁-C₆ alkoxy, fluoro, C(O)OR₁₁, aryl C₁-C₃ alkoxy, substituted aryl C₁-C₃ alkoxy, and the aryl substituents are as defined for R₃, R₄ and R₅;

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof.

Preferred substituents when R₂₀ and R₂₁ are taken together
5 include 7-aza-bicyclo[2.2.1]heptane and 2-aza-bicyclo[2.2.2]octane.

Unless otherwise stated or indicated, the following definitions shall apply throughout the specification and claims.

When any variable (e.g., aryl, heterocycle, R₁, etc.) occurs
10 more than one time in any constituent or in formula I, its definition on each occurrence is independent of its definition at every other occurrence. Also, combinations of substituents and/or variables are permissible only if such combinations result in stable compounds.

The term "alkyl" is intended to include both branched-
15 and straight-chain saturated aliphatic hydrocarbon groups having the specified number of carbon atoms, e.g., methyl (Me), ethyl (Et), propyl, butyl, pentyl, hexyl, heptyl, octyl, nonanyl, decyl, undecyl, dodecyl, and the isomers thereof such as isopropyl (i-Pr), isobutyl (i-Bu), sec-butyl (s-Bu), tert-butyl (t-Bu), isopentane, isohexane, etc.

20 The term "aryl" includes phenyl and naphthyl. Preferably, aryl is phenyl.

The term "halogen" or "halo" is intended to include fluorine, chlorine, bromine and iodine.

The term "heterocycle" or "heterocyclic ring" is defined
25 by all non-aromatic, heterocyclic rings of 3-7 atoms containing 1-3 heteroatoms selected from N, O, and S, such as oxirane, oxetane, tetrahydrofuran, tetrahydropyran, pyrrolidine, piperidine, tetrahydropyridine, tetrahydropyrimidine, tetrahydrothiophene, tetrahydrothiopyran, morpholine, hydantoin, valerolactam,
30 pyrrolidinone, and the like.

As used herein, the term "composition" is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified
35 amounts.

In addition, it is well known to those skilled in the art that many of the foregoing heterocyclic groups can exist in more than one tautomeric form. It is intended that all such tautomers be included within the ambit of this invention.

5 The optical isomeric forms, that is mixtures of enantiomers, e.g., racemates, or diastereomers as well as individual enantiomers or diastereomers of the instant compound are included. These individual enantiomers are commonly designated according to the optical rotation they effect by the symbols (+) and (-), (L) and (D), (1)
10 and (d) or combinations thereof. These isomers may also be designated according to their absolute spatial configuration by (S) and (R), which stands for sinister and rectus, respectively.

 The individual optical isomers may be prepared using conventional resolution procedures, e.g., treatment with an appropriate
15 optically active acid, separating the diastereomers and then recovering the desired isomer. In addition, the individual optical isomers may be prepared by asymmetric synthesis.

 Additionally, a given chemical formula or name shall encompass pharmaceutically acceptable addition salts thereof and
20 solvates thereof, such as hydrates.

 The compounds of the present invention, while effective themselves, may be formulated and administered in the form of their pharmaceutically acceptable addition salts for purposes of stability, convenience of crystallization, increased solubility and other desirable
25 properties.

 The compounds of the present invention may be administered in the form of pharmaceutically acceptable salts. The term "pharmaceutically acceptable salt" is intended to include all acceptable salts. Examples of acid salts are hydrochloric, nitric,
30 sulfuric, phosphoric, formic, acetic, trifluoroacetic, propionic, maleic, succinic, malonic, methane sulfonic and the like which can be used as a dosage form for modifying the solubility or hydrolysis characteristics or can be used in sustained release or prodrug formulations. Depending on the particular functionality of the compound of the present invention,
35 pharmaceutically acceptable salts of the compounds of this invention


include those formed from cations such as sodium, potassium, aluminum, calcium, lithium, magnesium, zinc, and from bases such as ammonia, ethylenediamine, N-methyl-glutamine, lysine, arginine, ornithine, choline, N,N'-dibenzylethylenediamine, chlorprocaine, 5 diethanolamine, procaine, N-benzylphenethylamine, diethylamine, piperazine, tris(hydroxymethyl)aminomethane, and tetramethyl-ammonium hydroxide. These salts may be prepared by standard procedures, e.g. by reacting a free acid with a suitable organic or inorganic base, or alternatively by reacting a free base with a suitable 10 organic or inorganic acid.


Also, in the case of an acid (-COOH) or alcohol group being present, pharmaceutically acceptable esters can be employed, e.g. methyl, ethyl, butyl, acetate, maleate, pivaloyloxymethyl, and the like, and those esters known in the art for modifying solubility or hydrolysis 15 characteristics for use as sustained release or prodrug formulations.

The compounds of the present invention may have chiral centers other than those centers whose stereochemistry is depicted in formula I, and therefore may occur as racemates, racemic mixtures and as individual enantiomers or diastereomers, with all such 20 isomeric forms being included in the present invention as well as mixtures thereof. Furthermore, some of the crystalline forms for compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds of the instant invention may form 25 solvates with water or common organic solvents. Such solvates are encompassed within the scope of this invention.

The compounds of the invention are prepared by the following reaction schemes. All substituents are as defined above unless indicated otherwise.

Scheme A






Reaction Scheme A

- As shown in reaction Scheme A, treatment of tryptamine
- 5 (1) with *N*-carboxyphthalimide in an inert organic solvent such as tetrahydrofuran at a temperature of 20-65°C, preferably 65°C, for a period of 12-48 hours gives the corresponding *N*-phthalimidotryptamine derivative (2). The *N*-phthalimidotryptamine (2) could be further
- 10 modified by treatment with a brominating agent such as pyridinium hydrobromide perbromide, pyrrolidone hydrotribromide, or the like in an inert organic solvent such as tetrahydrofuran, methylene chloride, chloroform, or mixtures thereof at 0-25°C for a period of 30 minutes to 4
- 15 hours to provide the 2-bromotryptamine (3). Bromide (3) may be reacted with an arylboronic acid (prepared essentially as described in : Gronowitz, S.; Hornfeldt, A.-B.; Yang, Y.-H. *Chem. Scr.* **1986**, 26, 311-314.) with palladium (0) catalysis, a weak base such as aqueous sodium carbonate or the like, and a chloride source such as lithium chloride in an inert solvent like toluene, benzene, ethanol, propanol or mixtures thereof at a temperature of 25°-100°C, preferably 80°C, for a period of 1-6
- 20 hours to give the 2-aryltryptamine derivative (4). Finally, the phthalimido group may be removed by treatment of (4) with aqueous hydrazine in an inert solvent such as methanol or ethanol at a temperature of 0°-25°C for a period of 4-24 hours to give tryptamine (5).

25

Scheme B



5


Reaction Scheme B

As shown in reaction Scheme B, the 2-aryltryptamine may be condensed with a carboxylic acid of type (6) using the coupling reagent 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), 1,3-dicyclohexylcarbodiimide (DCC) or the like with or without 1-hydroxybenzotriazole (HOBT) and a tertiary amine base such as N-methylmorpholine (NMM), triethylamine or the like in an inert organic solvent such as methylene chloride, chloroform, dimethylformamide, or mixtures thereof at or near room temperature for a period of 3-24 hours to provide the corresponding amide derivative (7). Alternatively, 2-aryltryptamine (5) can be treated with an active ester or acid chloride of type (8) in an inert organic solvent such as methylene chloride, chloroform, tetrahydrofuran, diethyl ether, or the like and a tertiary

amine base such as triethylamine, diisopropylethylamine, pyridine or the like at a temperature of 0°-25°C for 30 minutes to 4 hours to give (7).

Scheme C


5

Reaction Scheme C

As shown in reaction Scheme C, the amide carbonyl of (7) can be reduced by treatment with borane, lithium aluminum hydride, or equivalent hydride sources in an inert organic solvent such as tetrahydrofuran, diethyl ether, 1,4-dioxane or the like at 25°-100°C, preferably 65°C, for a period of 1-8 hours to give the corresponding amine compound (9).

15

Scheme D




5

Reaction Scheme D

As shown in reaction Scheme D, the 2-aryltryptamine (5) can be modified by treatment with an aldehyde or ketone of type (10) in the presence of a weak acid such as trifluoroacetic acid (TFA), acetic acid or the like, with or without a dessicant such as 3Å molecular sieves or magnesium sulfate, and a hydride source such as sodium borohydride or sodium cyanoborohydride, in an inert organic solvent such as methanol, ethanol, isopropanol, tetrahydrofuran, dichloromethane, chloroform, or mixtures thereof at a temperature of 0°-25°C for a period of 1-12 hours to give the corresponding secondary or tertiary amine derivative (11).

Scheme E




5

Reaction Scheme E

As shown in reaction Scheme E, treatment of an arylhydrazine or arylhydrazine hydrochloride (12) with an arylcyclopropylketone of type (13) in a polar organic solvent such as methanol, ethanol, n-propanol, isopropanol, n-butanol, t-butanol, preferably n-butanol, at a temperature of 70°-120°C for a period of 8-24 hours gives 2-aryltryptamine (5). Alternatively, when an arylhydrazine or arylhydrazine hydrochloride (12) is treated with an arylbutyl ketone of type (14) containing a leaving group (chloride, bromide, iodide, *O*-methansulfonate, *O*-trifluoromethansulfonate, or the like) at the 4-position in a polar solvent such as methanol, ethanol, n-propanol, isopropanol, n-butanol, t-butanol, or mixtures thereof at room temperature for a period of 30 minutes to 2 hours followed by heating to a temperature of 65°-100°C for 4-24 hours, 2-aryltryptamine (5) is produced.

20

Scheme F




5

Reaction Scheme F

As shown in reaction Scheme F, iodoanilines of type (15) may be reacted with aryl acetylenes, an appropriate palladium (0) catalyst such as tetrakis(triphenylphosphine)palladium, a copper (I) halide such as cuprous bromide in an inert organic solvent such as triethylamine at a temperature of 50°-88°C for a period of 30 minutes to 5 hours to provide the diacyl compound (16). Acetylene (16) may be further modified by treatment with a palladium (II) catalyst such as palladium (II) chloride or palladium (II) acetate in an inert organic solvent such as acetonitrile at a temperature of 50°- 82°C for a period of 30 minutes to 6 hours to give 2-arylidole (17).

Scheme G



5

Reaction Scheme G

As shown in reaction Scheme G, treatment of 2-arylimidole (17) with oxalyl chloride neat or in an inert organic solvent such as methylene chloride, chloroform, dichloroethane, tetrahydrofuran or the like at a temperature of 25°-65°C for a period of 3-24 hours gives the acylchloride adduct (18). The crude product (18) may be reacted with an amine of type (19) in an inert organic solvent such as diethylether, tetrahydrofuran, methylene chloride, chloroform or the like and an amine base such as triethylamine, diisopropylethylamine or pyridine at a temperature of 0°C-25°C for a period of 30 minutes to 4 hours to provide the amide derivative (20). Amide (20) may be further modified by


10

15

treatment with a reducing agent such as borane or lithium aluminum hydride in an inert organic solvent such as tetrahydrofuran at elevated temperatures, preferably reflux, for a period of 1-5 hours to give compound (21).

5

Scheme H



10


Reaction Scheme H

As shown in reaction Scheme H, N-benzyl derivatives of type (22a) or N-benzyloxycarbonyl derivatives of type (22b) may be reduced to provide the secondary amine analogs (7) by treatment with hydrogen (1 atm) and an appropriate catalyst such as palladium on carbon,


15

palladium hydroxide on carbon, or the like in an inert organic solvent such as tetrahydrofuran, ethyl acetate, methanol, ethanol, or mixtures thereof to which has been added a weak acid such as 30% aqueous acetic acid for a period of 10 minutes to 3 hours or until the aryl group has been removed to give the secondary amine.

Scheme I




10

Reaction Scheme I

As shown in reaction Scheme I, treatment of a nitroindole of type (24) with hydrogen (1 atm) and an appropriate catalyst such as Raney® Nickel in an inert organic solvent such as ethanol, methanol, or the like at room temperature for a period of 2-12 hours gives the corresponding aminoindole derivative (25).

Scheme J



5


Reaction Scheme J

As shown in reaction Scheme J, amino- or hydroxyindole (25) may be modified by acylation under a variety of conditions. For example, treatment of (25) with an acid chloride, acid anhydride or active ester and an amine base such as triethylamine, diisopropylethylamine, pyridine, or the like in an inert organic solvent such as methylene chloride, chloroform, tetrahydrofuran, or mixtures thereof at 0°C to room temperature for a period of 1 to 12 hours gives the corresponding amide or ester derivatives (26). Alternatively (25) may be coupled with a carboxylic acid by one of the many dehydrating agents commonly employed. For instance, treatment of aminoindole (25) with an appropriate carboxylic acid and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), 1,3-dicyclohexylcarbodiimide (DCC) or the like with or without 1-hydroxybenzotriazole (HOBT) and a tertiary amine base such as N-methylmorpholine (NMM), triethylamine or the like in an inert organic solvent such as methylene chloride,

chloroform, dimethylformamide, or mixtures thereof at or near room temperature for a period of 3-24 hours provides the corresponding amide or ester derivative (26).

5

Scheme K




Reaction Scheme K

- As shown in reaction Scheme K, urea or carbamate derivatives of (25) can be prepared by treatment with a carbamoyl chloride of type (27a), or alternatively with an isocyanate reagent of type (27b), and an amine base such as pyridine, triethylamine, diisopropylethylamine, *N*-methylmorpholine or the like in an inert organic solvent such as methylene chloride, chloroform, dimethylformamide, tetrahydrofuran or mixtures thereof at a temperature of 0°-65°C for a period of 1-72 hours to give (28). Compound (25) can also be modified by treatment with a bis(electrophilic) reagent such as phosgene, triphosgene, 1,1'-carbonyldiimidazole, *N,N'*-disuccinimidyl carbonate, or the like with or without the addition of an amine base such as pyridine, triethylamine, diisopropylethylamine, *N*-methylmorpholine in an inert solvent such as methylene chloride,

chloroform, or the like at a temperature of -20° - 0°C for a period of 20 minutes to 2 hours. After this time, the reaction mixture is treated with an appropriate mono- or disubstituted amine at -20° to 25°C for a period of 1-5 hours to give the urea or carbamate analog (28).

5


Scheme L



Reaction Scheme L

As shown in reaction Scheme L, amine (25) can be modified by treatment with an appropriate sulfonyl chloride of type (29) or sulfamyl chloride of type (30) with an amine base such as pyridine, triethylamine, diisopropylethylamine, *N*-methylmorpholine in an inert solvent such as methylene chloride, chloroform, dichloroethane or the like at a temperature of -20°-25°C for a period of 20 minutes to 2 hours to give the corresponding *N*-sulfonamide (31) or *N*-sulfamylamide (32) derivatives, respectively.


Scheme M



Reaction Scheme M

As shown in reaction Scheme M, the 2-aryltryptamine (33) can be modified by treatment with an epoxide such as (34) in an inert organic solvent such as methanol, ethanol, isopropanol, butanol, tert-butanol, or mixtures thereof at a temperature of 65°-110°C for a period of 8-20 hours to give the corresponding amino-alcohol derivative (35).


Scheme N



Reaction Scheme N

As shown in reaction Scheme N, amide derivatives of an acid-containing indole derivative such as (36) can be prepared by treatment with an appropriate amine ($R_{12}R_{11}NH$) and a suitable coupling agent such as benzotriazol-1-yloxy-tris(pyrrolidino)phosphonium hexafluorophosphate (PyBOP), benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate (BOP), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC), 1,3-dicyclohexylcarbodiimide (DCC) or the like with or without 1-hydroxybenzotriazole (HOBt) and a tertiary amine base such as N-methylmorpholine (NMM), triethylamine or the like in an inert organic solvent such as methylene chloride, chloroform, tetrahydrofuran, dimethylformamide, or mixtures thereof at or near room temperature for a period of 3-24 hours provides the corresponding amide derivative (37).


Scheme O



Reaction Scheme O

- 5 As shown in reaction Scheme O, pyridine derivatives such
as (38) can be oxidized to the corresponding N-oxide by treatment with an
oxidant such as 30% aqueous hydrogen peroxide and 0.5-5 mol% of a
catalyst such as methyltrioxorhenium(VII) or the like in an inert
organic solvent such as methylene chloride at or near room temperature
10 for 1-3 days to provide (39).

Scheme P



5

Reaction Scheme P

As shown in reaction Scheme P, the tryptamine 5 can be modified by reaction with an arylsulfonyl chloride such as 2-nitrobenzenesulfonyl chloride, 4-nitrobenzenesulfonyl chloride or 2,4-dinitrobenzenesulfonyl chloride and a hindered amine base such as 2,4,6-collidine, 2,6-lutidine or the like in an inert organic solvent such as methylene chloride to provide the corresponding sulfonamide 41. Sulfonamides such as 41 can be further modified by reaction with an alcohol of type 40 in the presence of triphenylphosphine and an activating agent such as diethyl azodicarboxylate (DEAD), diisopropyl

azodicarboxylate or the like in an inert organic solvent such as benzene, toluene, tetrahydrofuran or mixtures thereof to give the dialkyl-sulfonamide adduct. Removal of the sulfonyl group is accomplished by treatment with a nucleophilic amine such as n-propylamine or the like
5 in an inert organic solvent such as methylene chloride to give secondary amines of type 23.

The compounds of the present invention are useful in the treatment of various sex-hormone related conditions in men and
10 women. This utility is manifested in their ability to act as antagonists of the neuropeptide hormone GnRH as demonstrated by activity in the following *in vitro* assays.

Human GnRH receptor binding assay

15 Crude membranes prepared from CHO cells expressing human GnRH receptors were the sources for GnRH receptor. [¹²⁵I]Buserelin (a peptidyl GnRH analog) was used as the radiolabelled ligand. The binding activity was determined as an IC₅₀ which is the antagonist concentration required to inhibit the specific binding of [¹²⁵I]buserelin
20 to GnRH receptors by 50%.

Rat pituitary GnRH receptor binding assay:

Crude plasma membranes prepared from rat pituitary tissues were incubated in a Tris.HCl buffer (50 mM, PH. 7.5) containing bovine
25 serum albumin (.1%), [I-125]D-t-Bu-Ser6-Pro9-ethyl amide-GnRH, and the desired concentration of a test compound. The assay mixtures were incubated at 4°C for 90-120 minutes followed by rapid filtration and repeated washings through a glass fiber filter. The radioactivity of membrane bound radioligands was determined in a gamma-counter.
30 From this data, the IC₅₀ of the radioligand binding to GnRH receptors in the presence of test compound was estimated.

Inhibition of LH release assay:

Active compounds from the GnRH receptor binding assay were further evaluated with an *in vitro* LH release assay to confirm their antagonist activity (blocking GnRH-induced LH release).

5 1. Sample Preparation

The compounds to be assayed were dissolved and diluted in DMSO. The final concentration of DMSO in the incubation medium was 0.5%.

2. Assay

- 10 The Wistar male rats (150-200 grams) were obtained from Charles River Laboratories (Wilmington, MA). Rats were maintained at a constant temperature (25°C) on a 12-hr light, 12-hr dark cycle. Rat chow and water were available *ad libitum*. The animals were sacrificed by
- 15 in Hank's Balanced Salt Solution (HBSS) in a 50-mL polypropylene centrifuge tube. The collection tube was centrifuged for 5 min at 250 x g, and HBSS was removed by aspiration. Pituitary glands were transferred to a disposable petri plate and minced with a scalpel. The minced tissue
- 20 was then transferred to a 50-mL disposable centrifuge tube by suspending the tissue fragments in three successive 10-mL aliquots of HBSS containing 0.2% collagenase and 0.2% hyaluronidase. The cell dispersion was carried out in a water bath at 37°C with gentle stirring for 30 min. At the end of the incubation, the cells were aspirated 20 to 30
- 25 times with a pipet and the undigested pituitary fragments were allowed to settle for 3 to 5 min. The suspended cells were removed by aspiration, and then subjected to a 1200 x g centrifugation for 5 min. The cells were then resuspended in Culture medium. The undigested pituitary
- 30 fragments were treated with 30 mL aliquots of the digestion enzymes as above for a total of 3 digestions with the collagenase/hyaluronidase mixture. The resulting cell suspensions were pooled, counted and
- 35 diluted to a concentration of 3×10^5 cells/ml, and 1.0 ml of this suspension was placed in each well of a 24-well tray (Costar, Cambridge, MA). Cells were maintained in a humidified 5% CO₂-95% air atmosphere at 37°C for 3 to 4 days. The culture medium consisted of DMEM containing 0.37% NaHCO₃, 10% horse serum, 2.5% fetal bovine

serum, 1% non-essential amino acids, 1% glutamine, and 0.1% gentamycin. On the day of an experiment, cells were washed three times 1 1/2 hrs prior to and two more times immediately before the start of the experiment with DMEM containing 0.37% NaHCO₃, 10% horse
5 serum, 2.5% fetal bovine serum, 1% non-essential amino acids(100X), 1% glutamine(100X), 1% Penicillin/Streptomycin(10,000 Units of Penicillin and 10,000 micrograms of Streptomycin per ml), and 25 mM HEPES, pH 7.4. LH release was initiated by adding 1 ml of fresh medium containing test compounds in the presence of 2 nM GnRH to
10 each well in duplicate. Incubation was carried out at 37°C for 3 hr. After incubation, medium was removed and centrifuged at 2,000 x g for 15 min to remove any cellular material. The supernatant fluid was removed and assayed for LH content with a double antibody RIA procedure using materials obtained from Dr. A. F. Parlow (Harbor-
15 UCLA Medical Center, Torrance, CA).

The compounds of formula I are useful in a number of areas affected by GnRH. They may be useful in sex-hormone related conditions, sex-hormone dependent cancers, benign prostatic hypertrophy or myoma of the uterus. Sex-hormone dependent cancers which
20 may benefit from the administration of the compounds of this invention include prostatic cancer, uterine cancer, breast cancer and pituitary gonadotrophe adenomas. Other sex-hormone dependent conditions which may benefit from the administration of the compounds of this invention include endometriosis, polycystic ovarian disease, uterine
25 fibroids and precocious puberty. The compounds may also be used in combination with an angiotensin-converting enzyme inhibitor such as Enalapril or Captopril, an angiotensin II-receptor antagonist such as Losartan or a renin inhibitor for the treatment of uterine fibroids.

The compounds of the invention may also be useful for
30 controlling pregnancy, as a contraceptive in both men and women, for *in vitro* fertilization, in the treatment of premenstrual syndrome, in the treatment of lupus erythematosus, in the treatment of hirsutism, in the treatment of irritable bowel syndrome and for the treatment of sleep disorders such as sleep apnea.

A further use of the compounds of this invention is as an adjunct to growth hormone therapy in growth hormone deficient children. The compounds may be administered with growth hormone or a compound which increases the endogenous production or release of growth hormone. Certain compounds have been developed which stimulate the release of endogenous growth hormone. Peptides which are known to stimulate the release of endogenous growth hormone include growth hormone releasing hormone, the growth hormone releasing peptides GHRP-6 and GHRP-1 (described in U.S. Patent No. 4,411,890, PCT Patent Pub. No. WO 89/07110, and PCT Patent Pub. No. WO 89/07111) and GHRP-2 (described in PCT Patent Pub. No. WO 93/04081), as well as hexarelin (J. Endocrinol Invest., 15(Suppl 4), 45 (1992)). Other compounds which stimulate the release of endogenous growth hormone are disclosed, for example, in the following: U.S. Patent No. 3,239,345; U.S. Patent No. 4,036,979; U.S. Patent No. 4,411,890; U.S. Patent No. 5,206,235; U.S. Patent No. 5,283,241; U.S. Patent No. 5,284,841; U.S. Patent No. 5,310,737; U.S. Patent No. 5,317,017; U.S. Patent No. 5,374,721; U.S. Patent No. 5,430,144; U.S. Patent No. 5,434,261; U.S. Patent No. 5,438,136; EPO Patent Pub. No. 0,144,230; EPO Patent Pub. No. 0,513,974; PCT Patent Pub. No. WO 94/07486; PCT Patent Pub. No. WO 94/08583; PCT Patent Pub. No. WO 94/11012; PCT Patent Pub. No. WO 94/13696; PCT Patent Pub. No. WO 94/19367; PCT Patent Pub. No. WO 95/03289; PCT Patent Pub. No. WO 95/03290; PCT Patent Pub. No. WO 95/09633; PCT Patent Pub. No. WO 95/11029; PCT Patent Pub. No. WO 95/12598; PCT Patent Pub. No. WO 95/13069; PCT Patent Pub. No. WO 95/14666; PCT Patent Pub. No. WO 95/16675; PCT Patent Pub. No. WO 95/16692; PCT Patent Pub. No. WO 95/17422; PCT Patent Pub. No. WO 95/17423; Science, 260, 1640-1643 (June 11, 1993); Ann. Rep. Med. Chem., 28, 177-186 (1993); Bioorg. Med. Chem. Ltrs., 4(22), 2709-2714 (1994); and Proc. Natl. Acad. Sci. USA 92, 7001-7005 (July 1995).

Representative preferred growth hormone secretagogues employed in the present combination include the following:

- 1) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 5 2) N-[1(R)-[(1,2-Dihydro-1-methanecarbonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 10 3) N-[1(R)-[(1,2-Dihydro-1-benzenesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 15 4) N-[1(R)-[(3,4-Dihydro-spiro[2H-1-benzopyran-2,4'-piperidin]-1'-yl)carbonyl]-2-(1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 20 5) N-[1(R)-[(2-Acetyl-1,2,3,4-tetrahydrospiro[isoquinolin-4,4'-piperidin]-1'-yl)carbonyl]-2-(indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 25 6) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethyloxy)ethyl]-2-amino-2-methylpropanamide;
- 30 7) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethyloxy)ethyl]-2-amino-2-methylpropanamide methanesulfonate;
- 8) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(2',6'-difluorophenylmethyloxy)ethyl]-2-amino-2-methylpropanamide;
- 9) N-[1(R)-[(1,2-Dihydro-1-methanesulfonyl-5-fluorospiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethyloxy)ethyl]-2-amino-2-methylpropanamide;

- 10) N-[1(S)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl) carbonyl]-2-(phenylmethylthio)ethyl]-2-amino-2-methylpropanamide;
- 11) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-3-phenylpropyl]-2-amino-2-methylpropanamide;
- 12) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-3-cyclohexylpropyl]-2-amino-2-methylpropanamide;
- 13) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-4-phenylbutyl]-2-amino-2-methylpropanamide;
- 14) N-[1(R)-[(1,2-Dihydro-1-methanesulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(5-fluoro-1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 15) N-[1(R)-[(1,2-Dihydro-1-methanesulfonyl-5-fluorospiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(5-fluoro-1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 16) N-[1(R)-[(1,2-Dihydro-1-(2-ethoxycarbonyl)methylsulfonylspiro[3H-indole-3,4'-piperidin]-1'-yl)carbonyl]-2-(1H-indol-3-yl)ethyl]-2-amino-2-methylpropanamide;
- 17) N-[1(R)-[(1,2-Dihydro-1,1-dioxospiro[3H-benzothiophene-3,4'-piperidin]-1'-yl)carbonyl]-2-(phenylmethoxy)ethyl]-2-amino-2-methylpropanamide;

and pharmaceutically acceptable salts thereof.

The compounds of the invention may also be used in combination with bisphosphonates (bisphosphonic acids) and other agents, such as growth hormone secretagogues, e.g. MK-0677, for the

5 treatment and the prevention of disturbances of calcium, phosphate and bone metabolism, in particular, for the prevention of bone loss during therapy with the GnRH antagonist, and in combination with estrogens, progesterones and or androgens for the prevention or treatment of bone loss or hypogonadal symptoms such as hot flashes during therapy with the GnRH antagonist.

Bisphosphonates (bisphosphonic acids) are known to inhibit bone resorption and are useful for the treatment of bone lithiasis as disclosed in U.S. Patent 4,621,077 to Rosini, et al.

10 The literature discloses a variety of bisphosphonic acids which are useful in the treatment and prevention of diseases involving bone resorption. Representative examples may be found in the following: U.S. Patent No. 3,251,907; U.S. Patent No. 3,422,137; U.S. Patent No. 3,584,125; U.S. Patent No. 3,940,436; U.S. Patent No. 3,944,599; 15 U.S. Patent No. 3,962,432; U.S. Patent No. 4,054,598; U.S. Patent No. 4,267,108; U.S. Patent No. 4,327,039; U.S. Patent No. 4,407,761; U.S. Patent No. 4,578,376; U.S. Patent No. 4,621,077; U.S. Patent No. 4,624,947; U.S. Patent No. 4,746,654; U.S. Patent No. 4,761,406; U.S. Patent No. 4,922,007; U.S. Patent No. 4,942,157; U.S. Patent No. 5,227,506; U.S. 20 Patent No. 5,270,365; EPO Patent Pub. No. 0,252,504; and J. Org. Chem., 36, 3843 (1971).

The preparation of bisphosphonic acids and halo-bisphosphonic acids is well known in the art. Representative examples may be found in the above mentioned references which disclose the 25 compounds as being useful for the treatment of disturbances of calcium or phosphate metabolism, in particular, as inhibitors of bone resorption.

Preferred bisphosphonates are selected from the group of the following compounds: alendronic acid, etidrononic acid, clodronic acid, pamidronic acid, tiludronic acid, risedronic acid, 30 6-amino-1-hydroxy-hexylidene-bisphosphonic acid, and 1-hydroxy-3(methylpentylamino)-propylidene-bisphosphonic acid; or any pharmaceutically acceptable salt thereof. A particularly preferred bisphosphonate is alendronic acid (alendronate), or a pharmaceutically acceptable salt thereof. An especially preferred 35 bisphosphonate is alendronate sodium, including alendronate sodium

trihydrate. Alendronate sodium has received regulatory approval for marketing in the United States under the trademark FOSAMAX®.

Additionally, a compound of the present invention may be co-administered with a 5a-reductase 2 inhibitor, such as finasteride
5 or epristeride; a 5a-reductase 1 inhibitor such as 4,7b-dimethyl-4-aza-5a-cholestan-3-one, 3-oxo-4-aza-4,7b-dimethyl-16b-(4-chlorophenoxy)-5a-androstane, and 3-oxo-4-aza-4,7b-dimethyl-16b-(phenoxy)-5a-androstane as disclosed in WO 93/23420 and WO 95/11254; dual
10 inhibitors of 5a-reductase 1 and 5a-reductase 2 such as 3-oxo-4-aza-17b-(2,5-trifluoromethylphenyl-carbamoyl)-5a-androstane as disclosed in WO 95/07927; antiandrogens such as flutamide, casodex and cyproterone acetate, and alpha-1 blockers such as prazosin, terazosin, doxazosin, tamsulosin, and alfuzosin.

Further, a compound of the present invention may be used
15 in combination with growth hormone, growth hormone releasing hormone or growth hormone secretagogues, to delay puberty in growth hormone deficient children, which will allow them to continue to gain height before fusion of the epiphyses and cessation of growth at puberty.

For combination treatment with more than one active
20 agent, where the active agents are in separate dosage formulations, the active agents may be administered separately or in conjunction. In addition, the administration of one element may be prior to, concurrent to, or subsequent to the administration of the other agent.

The pharmaceutical compositions containing the active
25 ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions, hard or soft capsules, or syrups or elixirs. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical
30 compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically
35 acceptable excipients which are suitable for the manufacture of tablets.

These excipients may be for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example starch, gelatin or
5 acacia, and lubricating agents, for example, magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl
10 monostearate or glyceryl distearate may be employed. They may also be coated by the technique described in the U.S. Patent 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

Formulations for oral use may also be presented as hard
15 gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil.

Aqueous suspensions contain the active material in
20 admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring
25 phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethylene-oxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids
30 and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one

or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose, saccharin or aspartame.

Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil,
5 sesame oil or coconut oil, or in mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an
10 anti-oxidant such as ascorbic acid.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting
15 agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

The pharmaceutical compositions of the invention may also be in the form of an oil-in-water emulsions. The oily phase may be a
20 vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring phosphatides, for example soy beans, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation
25 products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavouring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such
30 formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable
35 dispersing or wetting agents and suspending agents which have been

mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water,
5 Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

10 Compounds of Formula I may also be administered in the form of a suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release
15 the drug. Such materials are cocoa butter and polyethylene glycols.

For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compound of Formula I are employed. (For purposes of this application, topical application shall include mouth washes and gargles.)

20 The compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal routes, using those forms of transdermal skin patches well known to those of ordinary skill in the art. To be administered in the form of a transdermal delivery system, the dosage
25 administration will, of course, be continuous rather than intermittent throughout the dosage regimen. Compounds of the present invention may also be delivered as a suppository employing bases such as cocoa butter, glycerinated gelatin, hydrogenated vegetable oils, mixtures of polyethylene glycols of various molecular weights and fatty acid esters
30 of polyethylene glycol.

The dosage regimen utilizing the compounds of the present invention is selected in accordance with a variety of factors including type, species, age, weight, sex and medical condition of the patient; the severity of the condition to be treated; the route of
35 administration; the renal and hepatic function of the patient; and the

particular compound thereof employed. A physician or veterinarian of ordinary skill can readily determine and prescribe the effective amount of the drug required to prevent, counter, arrest or reverse the progress of the condition. Optimal precision in achieving
5 concentration of drug within the range that yields efficacy without toxicity requires a regimen based on the kinetics of the drug's availability to target sites. This involves a consideration of the distribution, equilibrium, and elimination of a drug. Preferably, doses of the compound of structural formula I useful in the method
10 of the present invention range from 0.01 to 1000 mg per adult human per day. Most preferably, dosages range from 0.1 to 500 mg/day. For oral administration, the compositions are preferably provided in the form of tablets containing 0.01 to 1000 milligrams of the active ingredient, particularly 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0,
15 25.0, 50.0, 100 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied at a dosage level of from about 0.0002 mg/kg to about 50 mg/kg of body weight per day. The range is more particularly from about 0.001 mg/kg to 1
20 mg/kg of body weight per day.


Advantageously, the active agent of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in dividend doses of two, three or four times daily.

25 The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration.

It will be understood, however, that the specific dose level
30 for any particular patient will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

The following examples illustrate the preparation of some of the compounds of the invention and are not to be construed as limiting the invention disclosed herein.

5

EXAMPLE 1

10

2-(3,5-dimethylphenyl)-3-[2-(4-pyridin-4-ylbutylamino)ethyl]-1H-indol-5-ylmorpholin-4-ylmethanone

15 Step 1A 3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indole-5-carboxylic acid ethyl ester

A mixture of 7.60 g (50 mmol) of 4-hydrazinobenzoic acid, 10.55 g (50 mmol) of 3-chloropropyl 3,5-dimethylphenyl ketone, and 200 mL of absolute ethanol was stirred under nitrogen and heated to reflux. After 12 hours, the mixture was cooled and filtered. The solid on the filter was washed with additional small volumes of ethanol. The filtrate was treated with 4 mL of concentrated sulfuric acid and stirred at reflux under nitrogen for 4 days. The cooled mixture was stirred in an ice bath as a solution of sodium ethoxide (21% w/w in ethanol) was added dropwise until the mixture was basic by pH paper. The mixture was filtered and concentrated *in vacuo* at 30 °C. The residue was partitioned between diethyl ether and water, with some saturated aqueous sodium chloride solution added to assist in separation of the layers. The aqueous

phase was washed with an additional 100 mL of ether. The combined organic extracts were dried over sodium sulfate, filtered, and concentrated *in vacuo*. The residual gum was purified by flash chromatography on silica gel (elution with 97:3:0.3 and then 95:5:0.5 methylene chloride:methanol:-ammonium hydroxide) to give the title compound (4.8g). 400 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (PB-NH₃/CI): m/e = 337 (M + H).

Step 1B 2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-4-yl)butylamino]ethyl]-1H-indole-5-carboxylic acid ethyl ester
To a dry flask were added 5.0 g (14.9 mmol) of 3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indole-5-carboxylic acid ethyl ester, 1.98 g (13.5 mmol) of 4-(pyridin-4-yl)butyraldehyde (diluted with 0.5 mL of CDCl₃), 8.12 g (67.7 mmol) of anhydrous magnesium sulfate, and a magnetic stirring bar. The flask was purged with nitrogen, cooled to -10 °C, and stirred as 11.5 mL of dry CDCl₃ was introduced gradually by syringe. The mixture was stirred under nitrogen for about 20 minutes. Next, the septum was removed, and 670 mg (17.6 mmol) of sodium borohydride was added rapidly. The septum was immediately replaced, and the system was again purged with nitrogen. The mixture was stirred under nitrogen at about -5 °C as 10 mL of dry methanol was added gradually by syringe. After a few minutes at this temperature, the reaction was removed from the cooling bath and partitioned between 80 mL of ethyl acetate and 100 mL of water. The organic layer was dried over sodium sulfate, filtered, and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (elution with a gradient of 4-9% methanol in methylene chloride; repeated using 5-15% methanol in methylene chloride) to give the title compound (3.19 g). 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (PB-NH₃/CI): m/e = 470.4 (M + H). An additional 1.91 g of less pure material was also isolated.

Step 1C 3-[2-[benzyloxycarbonyl]-4-(pyridin-4-yl)butylamino]ethyl]-2-(3,5-dimethylphenyl)-1H-indole-5-carboxylic acid ethyl ester

A solution of 3.19 g (6.83 mmol) of 2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-4-yl)butylamino] ethyl]-1*H*-indole-5-carboxylic acid ethyl ester in 25 mL of dry methylene chloride was stirred under nitrogen and cooled to -78 °C in a dry ice-acetone bath as 2.38 mL (1.76 g, 13.7 mmol) of *N,N*-diisopropylethylamine was added, followed by gradual addition of 3.4 mL (4.06 g; 23.7 mmol) of benzyl chloroformate by syringe, in portions. After about 2.5 hours, the solution was removed from the cooling bath and allowed to warm to room temperature. It was then partitioned between ethyl acetate and 5% aqueous potassium bisulfate solution. The organic phase was dried over magnesium sulfate, filtered, and concentrated *in vacuo*. Purification of the residue by flash chromatography on silica gel (elution with a gradient of 0.5-10% methanol in methylene chloride) afforded a quantitative yield of the product as a yellow foam. 500 MHz ¹H NMR was complex, owing to the existence of rotamers, but was consistent with the assigned structure. Mass spectrum (PB-NH₃/CI): *m/e* = 604.3 (M + H).

Step 1D 3-[2-[benzyloxycarbonyl]-[4-(pyridin-4-yl)butyl]amino]-ethyl-2-(3,5-dimethylphenyl)-1*H*-indole-5-carboxylic acid hydrochloride

A solution of 4.11 g (6.83 mmol) of 3-[2-[benzyloxycarbonyl]-[4-(pyridin-4-yl)butyl]amino]ethyl-2-(3,5-dimethylphenyl)-1*H*-indole-5-carboxylic acid ethyl ester in 161 mL (80.5 mmol) of 0.50 N KOH in methanol was stirred at about 60 °C as 19 mL of water was added gradually. Stirring was continued at reflux overnight. The cooled mixture was concentrated *in vacuo* to give a yellow solid, which was partitioned between 250 mL of a 1:1 ethyl acetate-tetrahydrofuran mixture and 250 mL of 0.5 N HCl. The organic phase was washed twice with 0.5 N HCl, then dried over magnesium sulfate, filtered, and concentrated *in vacuo*. The resulting solid was triturated with diethyl ether and collected on a filter to give (after drying) 3.46 g of yellow solid, mp 133.5-137.5 °C; homogeneous by TLC (95:5:0.5 CH₂Cl₂-MeOH-AcOH). 500 MHz ¹H NMR (DMSO-*d*₆) was consistent with the assigned structure. Mass spectrum (ESI): *m/e* = 576.4 (M + H)⁺

Step 1E [2-[2-(3,5-dimethylphenyl)-5-(morpholine-4-carbonyl)-1H-indol-3-yl]ethyl]-(4-pyridin-4-ylbutyl)carbamic acid benzyl ester

A mixture of 100 mg (0.163 mmol) of 3-[2-[benzyloxycarbonyl-[4-(pyridin-4-yl)butyl]amino] ethyl]-2-(3,5-dimethylphenyl)-1H-indole-5-carboxylic acid hydrochloride 101.7 mg (0.196 mmol) of PyBOP reagent, 0.085 mL (85.2 mg, 0.978 mmol) of morpholine, and 1 mL of dry methylene chloride was stirred under nitrogen at room temperature in a stoppered flask. After 5 days, the solution was partitioned between ethyl acetate and saturated aqueous sodium bicarbonate solution. The organic phase was dried over sodium sulfate, filtered, and concentrated *in vacuo*. Purification of the crude product by flash chromatography on silica gel (gradient elution with 1-4% MeOH in CH₂Cl₂) afforded a quantitative yield of the title compound as a yellow gum; homogeneous by TLC in 95:5 CH₂Cl₂-MeOH. 500 MHz ¹H NMR (CDCl₃) was complex, owing to rotamers, but was consistent with the assigned structure. Mass spectrum (ESI): m/e = 645.6 (M + H).

Step 1F [2-(3,5-dimethylphenyl)-3-[2-(4-pyridin-4-ylbutylamino)-ethyl]-1H-indol-5-yl]morpholin-4-ylmethanone

A mixture of 113 mg (0.175 mmol) of [2-[2-(3,5-dimethylphenyl)-5-(morpholine-4-carbonyl)-1H-indol-3-yl]ethyl]-(4-pyridin-4-ylbutyl)carbamic acid benzyl ester, 50 mg of 20% palladium hydroxide on carbon, and 10 mL of 2-methoxyethanol was shaken with hydrogen (approx. 50 psig) in a pressure vessel for 2.5 hours. The catalyst was removed by filtration through Celite, and the filtrate was concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (gradient elution from 99:1:0.1 to 94:6:0.6 CH₂Cl₂-MeOH-concd. NH₄OH) yielded 53.2 mg (60%) of a white, stiff foam; homogeneous by TLC in 95:5:0.5 CH₂Cl₂-MeOH-concd. NH₄OH. 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (ESI): m/e = 511.5 (M + H).

PREPARATION OF SYNTHETIC INTERMEDIATES

Step A: 4-(4-pyridyl)-3-pentyn-1-ol

4-Bromopyridine hydrochloride salt (5.5 g) was dissolved in a solvent mixture comprising triethylamine (50 mL) and water (10 mL). Anhydrous lithium chloride (100 mg), copper (I) bromide powder (100 mg) and but-3-yn-1-ol (2.17g) was added to the pyridine salt and the mixture stirred as an active nitrogen gas stream passed gently through the solution for approximately 15 minutes after which time tetrakis(triphenylphosphine)palladium (250 mg) was added. The reaction mixture was heated to reflux under a nitrogen atmosphere and maintained at reflux for 2.5h after which heating was stopped and the reaction allowed to cool to room temperature. The mixture was concentrated *in vacuo* and the residue treated with 3 M sodium hydroxide, extracted with chloroform and concentrated *in vacuo*. Purification of the residue by flash chromatography on silica gel (ethyl acetate) gave the title compound (3.74 g).

Step B: 4-(4-pyridyl)-butan-1-ol

4-(4-Pyridyl)-3-butyne-1-ol (3.5g) was dissolved in methanol (100 mL) in a Parr hydrogenation bottle and platinum (IV) oxide [Adams' Catalyst] (0.3g) was added. The Parr bottle was placed on a Parr hydrogenation apparatus and the solution hydrogenated at 40 psi for 2.5h after which time the starting material had been judged to be consumed by TLC. The spent catalyst was removed by filtration through a Celite pad and the pad carefully washed with more methanol. The combined filtrates were evaporated under reduced pressure on a rotary evaporator and the oily residues then subjected to column chromatography on a short silica column using neat ethyl acetate as the eluant to provide the title compound (3.0 g).


Step C: 4-(pyridin-4-yl)butyraldehyde

Oxalyl chloride (1.45 mL of a 2M solution in dry methylene chloride) was placed in an oven-dried flask and cooled to -78°C using a dry ice and acetone cooling bath and a solution of DMSO (0.413 mL) in dry methylene chloride (1 mL) added drop by drop to the oxalyl chloride over 3 minutes and stirred for a further 3 minutes. A solution of 4-(4-

pyridyl)-butan-1-ol (400 mg) in dry methylene chloride (5 mL) was added to the reaction flask over approx. 3 minutes and the reaction stirred for 15 minutes. Anhydrous triethylamine (2.03 mL) was added and the reaction mixture stirred for another 2 hours during which time the cold
 5 bath had warmed up to room temperature. The reaction was quenched by the addition of saturated brine and then partitioned with methylene chloride. The aqueous layer was discarded and the methylene chloride extract dried over anhydrous sodium sulfate powder, filtered and evaporated under reduced pressure to leave an oily residue. The product
 10 was isolated by column chromatography on silica gel using ethyl acetate as eluant (301 mg).

EXAMPLE 2

15



20

1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[4-(4-pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropan-1-one dihydrochloride

Step 2A 2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropionic acid ethyl ester

25

A dry flask containing 3.00 g (7.93 mmol) of 2-[3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid ethyl ester (prepared essentially as described in EXAMPLE 1 from ethyl 2-(4-hydrazinophenyl)-2-methylpropionate), 4.76 g (39.7 mmol) of

anhydrous magnesium sulfate, and a magnetic stirring bar was fitted with a septum and needle adapter leading to a Firestone valve. The flask was thoroughly purged with nitrogen, and the mixture was cooled in an ice-methanol bath at -10 to -5 °C and stirred vigorously as a solution of 1.32 g (8.88 mmol) of 4-(pyridin-3-yl)butyraldehyde in 15 mL of dry CDCl_3 was added gradually by syringe over 10-15 minutes. The resulting mixture was stirred under nitrogen at -10 to -5 °C for 40-45 minutes. Then the septum was removed just long enough to add 390 mg (10.3 mmol) of sodium borohydride. The mixture was stirred under nitrogen at -10 to -5 °C as 10 mL of dry methanol was added dropwise by syringe over several minutes. After 30 minutes, the mixture was removed from the cooling bath and partitioned between 90 mL of ethyl acetate and 90 mL of water. The organic layer was washed with 2 x 30 mL of brine, then dried over anhydrous sodium sulfate. The filtered solution was concentrated *in vacuo*, and the residue was flash chromatographed of silica gel (gradient elution with 0-10% methanol in methylene chloride). Fractions containing product and a small amount of unreacted starting material were combined and concentrated to give 3.00 g of light beige, stiff foam, used directly in the next step without further purification or characterization.

Step 2B 2-[3-[2-[benzyloxycarbonyl]-4-(pyridin-3-yl)butyl]aminoethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid ethyl ester

A solution of 3.00 g (max. 5.86 mmol) of crude 2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropionic acid ethyl ester in 30 mL of dry methylene chloride was stirred under nitrogen with cooling in a dry-ice-acetone bath. To this solution was added by syringe 1.106 mL (820 mg, 6.36 mmol) of *N,N*-diisopropylethylamine. Then 0.956 mL (1.14 g, 6.36 mmol) of benzyl chloroformate was added dropwise by syringe over 5-10 minutes. After 20 minutes, the solution was removed from the cooling bath and allowed to warm to room temperature. After 2 hours, the solution was diluted with 50 mL of methylene chloride, transferred to a separatory funnel, and shaken with 80 mL of water. The organic phase was dried over

magnesium sulfate, filtered, and concentrated *in vacuo*. Flash chromatography of the residual gum on silica gel (gradient elution with 0.2-2% methanol in methylene chloride gave 2.81 g (55% overall for Steps 1 and 2) of pale, golden-yellow gum; virtually homogeneous by TLC in 95:5 CH₂Cl₂-MeOH. 500 MHz ¹H NMR (CDCl₃) was complex, owing to rotamers, but appeared to be consistent with the assigned structure. Mass spectrum (ESI): *m/e* = 646 (M + H).

Step 2C 2-[3-[2-[benzyloxycarbonyl]-[4-(pyridin-3-yl)butyl]amino]ethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-methylpropionic acid
A mixture of 2.78 g (4.30 mmol) of 2-[3-[2-[benzyloxycarbonyl]-[4-(pyridin-3-yl)butyl]amino]ethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-2-methylpropionic acid ethyl ester in 43.0 mol (21.5 mmol) of 0.5 M potassium hydroxide in methanol and 25 mL of tetrahydrofuran was stirred under nitrogen and heated to reflux. To the resulting solution was gradually added 18 mL of water, and the solution was maintained at reflux for 39 hours. It was then cooled and concentrated to small volume, accompanied by precipitation. The mixture was treated with 10.75 mL (21.5 mmol) of 2 N hydrochloric acid and agitated for a few minutes. The solid was collected on a filter and washed thoroughly with water. After suction-drying under nitrogen, the solid was triturated and washed with diethyl ether and vacuum-dried to yield 2.43 g (92%) of cream-colored powder, mp 152-154 °C (partial dec.); homogeneous by TLC in 90:10 CH₂Cl₂-MeOH. 500 MHz ¹H NMR (DMSO-*d*₆) was consistent with the assigned structure. Mass spectrum (ESI): *m/e* = 618 (M + H).

Step 2D [2-[5-[2-(7-azabicyclo[2.2.1]hept-7-yl)-1,1-dimethyl-2-oxoethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl]-[4-(pyridin-3-yl)butyl]carbamic acid benzyl ester
A mixture of 92.7 mg (0.15 mmol) of 2-[3-[2-[benzyloxycarbonyl]-[4-(pyridin-3-yl)butyl]amino]ethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-2-methylpropionic acid, 80.2 mg (0.6

mmol) of 7-azabicyclo[2.2.1]heptane hydrochloride, 83.2 mg (0.16 mmol) of PyBOP reagent, 0.107 mL (77.8 mg; 0.77 mmol) of triethylamine, and 0.75 mL of dry methylene chloride was stirred at room temperature in a stoppered flask for 48 hours. The solution was then partitioned between 5 10 mL of ethyl acetate and 10 mL of 0.5 N hydrochloric acid. The organic phase was washed with 10 mL of saturated aqueous sodium bicarbonate solution and then with 5 mL of saturated aqueous sodium chloride solution. The ethyl acetate phase was then dried (magnesium sulfate), filtered, and concentrated *in vacuo* at room temperature. The residue 10 was purified by preparative TLC on 6 Analtech tapered silica gel plates (20 x 20 cm), which were developed in 95:5 CH₂Cl₂-MeOH. The product band from each plate was isolated, combined, and extracted with 95:5 CH₂Cl₂-MeOH. Concentration of the extracts *in vacuo* yielded 85.9 mg (82%) of a very pale yellow glass; virtually homogeneous by TLC in 95:5 15 CH₂Cl₂-MeOH. 500 MHz ¹H NMR (CDCl₃) was complex, owing to rotamers, but was consistent with the assigned structure. Mass spectrum (ESI): *m/e* = 697.6 (M + H).

Step 2E 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylaminoethyl]-1H-indol-5-yl]-2-methylpropan-1-one

A mixture of 80.2 mg (0.115 mmol) of [2-[5-[2-(7-azabicyclo[2.2.1]hept-7-yl)-1,1-dimethyl-2-oxoethyl]-2-(3,5-dimethylphenyl)-1H-indol-3-yl]ethyl]-[4-(pyridin-3-yl)butyl]carbamic acid 25 benzyl ester, 40 mg of 10% palladium on carbon, 4 mL of absolute ethanol, and 4 mL of ethyl acetate was shaken with hydrogen (46 psig) in a pressure vessel for 6 hours. The catalyst was removed by filtration through Celite under nitrogen, and the filtrate was concentrated *in vacuo* at room temperature. The residue was purified by preparative 30 TLC on 4 Analtech tapered silica gel plates (20 x 20 cm), which were developed in 92.5:7.5:0.75 CH₂Cl₂-MeOH-concentrated NH₄OH. The product band from each plate was isolated, combined, and extracted with 92.5:7.5:0.75 CH₂Cl₂-MeOH-concentrated NH₄OH. Concentration of the extracts *in vacuo* yielded 53.8 mg (81%) of a pale yellow, stiff gum 35 or glass; essentially homogeneous by TLC in 92.5:7.5:0.75 CH₂Cl₂-

MeOH-concentrated NH_4OH . 500 MHz ^1H NMR (CDCl_3) was consistent with the assigned structure. Mass spectrum (ESI): $m/e = 563.5$ ($M + H$).

Step 2F 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropan-1-one dihydrochloride

5
A solution of 42.8 mg (0.0760 mmol) of 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropan-1-one in 1.5 mL of
10 methanol was treated with 0.152 mL (0.304 mmol) of 2 N hydrochloric acid. The solution was agitated and allowed to stand briefly before being filtered. The filtrate was evaporated to dryness under nitrogen, and the residue was triturated with diethyl ether. The resulting solid was collected on a filter, washed with additional ether, and dried to yield 46.5
15 mg (96%) of light golden-tan powder, mp $>160^\circ\text{C}$ (gradual; preliminary softening). 500 MHz ^1H NMR ($\text{DMSO}-d_6$) was consistent with the assigned structure.

20 PREPARATION OF SYNTHETIC INTERMEDIATES

Step A: Ethyl (+/-)-2-(4-nitrophenyl)propionate

To a solution of 9.76 g (50 mmol) of (+/-)-2-(4-nitrophenyl)propionic acid in 150 mL of absolute ethanol was added 3.0
25 mL of concentrated sulfuric acid. The resulting solution was stirred at reflux under nitrogen. After 6 hours, the solution was cooled and stirred vigorously as 250 mL of saturated aqueous sodium bicarbonate solution was added gradually (Caution: foaming). The mixture was then partitioned between 750 mL of ethyl acetate and 500 mL of water.
30 The organic layer was washed with 100 mL of saturated aqueous sodium bicarbonate solution and then with 100 mL of saturated aqueous sodium chloride solution. The organic phase was dried over magnesium sulfate, filtered, and concentrated *in vacuo* to give 10.86 g (97%) of an oil;

homogeneous by TLC in 9:1 hexane-ethyl acetate. 400 MHz ^1H NMR (CDCl_3) was consistent with the assigned structure.

Step B: Ethyl 2-methyl-2-(4-nitrophenyl)propionate

- 5 A suspension of 924 (23 mmol) of sodium hydride (60% in oil) in 21 mL of dry *N,N*-dimethylformamide was stirred under nitrogen in an ice bath as a solution of 4.68 g (21 mmol) of ethyl (+/-)-2-(4-nitrophenyl)propionate in 20.5 mL of dry *N,N*-dimethylformamide was added gradually over about 10 minutes. An intense violet color developed
- 10 during the addition. The mixture was then allowed to warm to room temperature. After about 1 hour, the mixture was again cooled in an ice bath as a solution of 1.44 mL (3.28 g; 23 mmol) of methyl iodide in 5 mL of dry *N,N*-dimethylformamide was added dropwise by syringe over about 10 minutes, while maintaining the internal temperature at 10-15 °C.
- 15 The mixture was allowed to warm to room temperature, and the color changed to brown. After 1 hour, an additional 187 mL (426 mg, 3 mmol) of iodomethane was added. By the next day, the mixture consisted of a suspension of some grayish solid in a golden liquid. It was stirred vigorously and quenched by gradual addition of 10 mL of 5% aqueous
- 20 potassium bisulfate solution. The mixture was partitioned between 400 mL of diethyl ether and 400 mL of water. The organic layer was washed with an additional 3 x 400 mL of water and then with 50 mL of saturated aqueous sodium chloride solution. The organic phase was then dried over magnesium sulfate, filtered, and concentrated *in vacuo*. Flash
- 25 chromatography of the residue on silica gel (elution with 19:1 hexane-ethyl acetate) yielded 4.31 g (87%) of an oil; homogeneous by TLC in 9:1 hexane-ethyl acetate. 400 MHz ^1H NMR (CDCl_3) was consistent with the assigned structure.

30 Step C: Ethyl 2-(4-Aminophenyl)-2-methylpropionate


A mixture of 4.27 g (18 mmol) of ethyl 2-methyl-2-(4-nitrophenyl)propionate, 200 mg of 10% palladium on carbon, and 120 mL of absolute ethanol was shaken with hydrogen (initial hydrogen pressure 47 psig) in a pressure vessel for 2 hours. The catalyst was

removed by filtration through Celite under nitrogen, and the filter cake was washed with additional ethanol. Concentration of the filtrate *in vacuo* at up to 50 °C gave 3.74 g (100%) of an oil; homogeneous by TLC in 4:1 hexane-EtOAc. 400 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (ESI): *m/e* = 208 (M + H).

Step D: Ethyl 2-(4-hydrazinophenyl)-2-methylpropionate

A solution of 3.725 g (18 mmol) of ethyl 2-(4-aminophenyl)-2-methylpropionate in 18 mL of concentrated hydrochloric acid was stirred at -10 to -5 °C in an ice-acetone bath as a solution of 1.29 g (18.7 mmol) of sodium nitrite in 7.5 mL of water was added dropwise over about 15 minutes. Stirring was continued at this temperature for an additional 30 minutes. Next, a small amount of insoluble solid was removed by filtration into a cold receiving flask. The filtrate was then added dropwise over 10-15 minutes to a solution of 20.3 g (90 mmol) of stannous chloride dihydrate in 14.5 mL of concentrated hydrochloric acid stirred under nitrogen in an ice-acetone bath. The addition was carried out at such a rate that the internal temperature remained at about -5 °C. A gummy material separated during the addition. After completion of the addition, stirring was continued at -10 to -5 °C for 1 hour. The aqueous phase was decanted, and the residual gum was dissolved in 250 mL of ethyl acetate. The ethyl acetate solution was treated cautiously with 250 mL of saturated aqueous sodium bicarbonate solution and shaken in a separatory funnel. The ethyl acetate layer was washed with 50 mL of saturated aqueous sodium chloride solution. The entire mixture was filtered before separation of the phases. The ethyl acetate phase was dried over magnesium sulfate, filtered, and concentrated *in vacuo* at room temperature to yield 2.59 g (65%) of an oil. 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure and indicated that only minor impurities were present.

EXAMPLE 3.1



5-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl]-1H-indol-5-yl]-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one

5

Step 3.1A 2-ethyl-4,4-dimethyl-5-(4-nitrophenyl)-2,4-dihydropyrazol-3-one

A mixture of 1.00 g (4 mmol) of 2,2-dimethyl-3-(4-nitrophenyl)-3-oxopropionic acid methyl ester (Yang, C.-Y.; Wnek, G. E., Polymer, **1992**, 33, 4191-4196), 3.00 g (20 mmol) of ethylhydrazine oxalate, 8 mL of 2-methoxyethanol, and 4 mL of glacial acetic acid was stirred under nitrogen at gentle reflux for 24 hours. The cooled solution was concentrated *in vacuo*. The residue was partitioned between ethyl acetate and water. The ethyl acetate phase was washed with additional water and then with saturated aqueous sodium chloride solution. The organic solution was dried over magnesium sulfate, filtered, and concentrated *in vacuo* to give 703 mg (67%) of light yellow crystals, mp 121-122 °C; homogeneous by TLC in 2:1 hexane-EtOAc. 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (PB-NH₃/CI): m/e = 232.1 (M - Et), 265.1 (M + H).

10

15

20

Step 3.1B 5-(4-aminophenyl)-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one

Hydrogenation of 2-ethyl-4,4-dimethyl-5-(4-nitrophenyl)-2,4-dihydropyrazol-3-one according to the procedure of Example 3.2 Step E afforded a quantitative yield of light yellow-tan solid, mp 118-120.5 °C; homogeneous by TLC in 1:1 hexane-EtOAc and 98:2 CH₂Cl₂-MeOH. 500

25

MHz ^1H NMR (CDCl_3) was consistent with the assigned structure. Mass spectrum (ESI): $m/e = 232.1$ ($M + H$).

5 Step 3.1C 2-ethyl-5-(4-hydrazinophenyl)-4,4-dimethyl-2,4-dihydropyrazol-3-one

10 This material was prepared from 5-(4-aminophenyl)-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one according to the procedure of Example 2 REACTION INTERMEDIATES, Step D, except that the entire reaction mixture from the stannous chloride reduction was stirred in an ice bath and treated cautiously with excess saturated sodium carbonate solution (CAUTION: foaming), resulting in precipitation. This material was transferred to a separatory funnel and shaken with 2:1 $\text{Et}_2\text{O}-\text{CH}_2\text{Cl}_2$. The mixture was filtered before separation of the phases. The aqueous phase was extracted further with several portions of ethyl acetate. The combined organic fractions were concentrated *in vacuo* to give an 80% yield of an amorphous, light yellow-orange solid, mp 131.5-135 °C dec; ill-defined by TLC in 95:5 CH_2Cl_2 -MeOH. 500 MHz ^1H NMR (CDCl_3) was consistent with the assigned structure.

20 Step 3.1D 5-[3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one


25 This compound was prepared from 2-ethyl-5-(4-hydrazinophenyl)-4,4-dimethyl-2,4-dihydropyrazol-3-one and 3-chloropropyl 3,5-dimethylphenyl ketone according to the procedure of Example 3.2 Step A, except that the reaction time was 15 hours. Flash chromatography of the crude product on silica gel (gradient elution with 97:3 and 95:5 CH_2Cl_2 -MeOH followed by 95:5:0.5 and 92.5:7.5:0.75 CH_2Cl_2 -MeOH-concd. NH_4OH) gave a 27% yield of light tan, stiff foam; homogeneous by TLC in 92.5:7.5:0.75 CH_2Cl_2 -MeOH-concd. NH_4OH . 500 MHz ^1H NMR (CDCl_3) was consistent with the assigned structure. Mass spectrum (PB- NH_3/CI): $m/e = 403.2$ ($M + H$).

Step 3.1E 5-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-3-yl)butylamino]ethyl-1H-indol-5-yl]-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one

A mixture of 82.5 mg (0.205 mmol) 5-[3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-ethyl-4,4-dimethyl-2,4-dihydropyrazol-3-one and 124 mg (0.475 mmol) of magnesium sulfate was purged with nitrogen and stirred in an ice-methanol bath at about -10 to -5 °C as a solution of 34.3 mg (0.23 mmol) of 4-(pyridin-4-yl)butyraldehyde in 0.500 mL of dry CDCl₃ was added gradually by syringe. The mixture was stirred under nitrogen at this temperature for 30 minutes. The septum was removed just long enough to add 10.0 mg (0.265 mmol) of sodium borohydride, and the solution was repurged with nitrogen. The mixture was stirred at -10 to -5 °C as 350 mL of dry methanol was added gradually, and stirring was continued at this temperature. After 45 minutes, the mixture was partitioned between ethyl acetate and water. The ethyl acetate layer was washed with brine, then dried over sodium sulfate, filtered, and concentrated *in vacuo*. The residue was purified by preparative TLC on 6 1000-micron silica gel GF plates (developed in 87.5:12.5 CH₂Cl₂-MeOH). Isolation of the product band (by extraction with 90:10:1 CH₂Cl₂-MeOH-concd. NH₄OH) gave 49.8 mg (45%) of a light beige, stiff foam; essentially homogeneous by TLC in 90:10 CH₂Cl₂-MeOH. 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum (ESI): m/e = 536.4 (M + H).

25

EXAMPLE 3.2



1-[2-(3,5-dimethylphenyl)-3-[2-(4-pyridin-3-yl-butylamino)ethyl]-1H-indol-5-yl]4-methyl-1,4-dihydrotetrazol-5-one

5 Step 3.2A 2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethylamine

To a solution of 3-chloropropyl 3,5-dimethylphenyl ketone (2.5g in 13.5 mL *tert*-butanol) was added 1.65 g 4-nitrophenylhydrazine and stirred for 20 minutes at room temperature. At this time, 108 mL of 90% aqueous methanol was added and the mixture heated to reflux on an oil bath. After 16 hours, the mixture was cooled to room temperature and the volatiles removed *in vacuo*. The residue was triturated with ethyl acetate and allowed to stand at 0 °C for 8 hours. Filtration of the resulting suspension gave the crude title compound as the hydrochloride salt (1.4 g).

15

Step 3.2 B N-[2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl]benzamide

To a solution of 2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethylamine (3.0 g in 80 mL dry methylene chloride) at 0 °C was added 4.0 mL triethylamine followed by 1.4 mL benzoyl chloride and the mixture stirred at low temperature. After 20 minutes, the reaction was quenched by the addition of saturated aqueous sodium bicarbonate and extracted with ethyl acetate. The organic portion was washed with water and concentrated *in vacuo* to give the crude title compound (1.43 g).

25

Step 3.2C Benzyl-[2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl]amine

To a stirred solution of N-[2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl]benzamide (1.7 g in 130 mL dry tetrahydrofuran) was added 35 mL of a 1M solution of borane in tetrahydrofuran and the mixture heated slowly to reflux on an oil bath. After 2 hours the mixture was cooled to room temperature and the excess borane quenched by the careful addition of methanol. The mixture was concentrated to half-volume, treated with *N,N*-

35

dimethylethanolamine (13 mL) and heated to reflux on an oil bath. After 3 hours the mixture was cooled to room temperature and concentrated *in vacuo*. Purification by flash chromatography on silica gel (methylene chloride:methanol, 97:3) gave the title compound (1.5 g).

5

Step 3.2D Benzyl-{2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl}-(4-pyridin-3-yl-butyl)amine

A mixture of benzyl-{2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl}amine (700 mg) and 4-pyridin-3-yl butyraldehyde (314 mg) were solvated in 30 mL dry methanol to which ca. 2 g powdered 3Å molecular sieves were added. The pH of this mixture was adjusted to 5 by the addition of trifluoroacetic acid and then 441 mg sodium cyanoborohydride was added and the mixture stirred at room temperature. After 48 hours, the mixture was filtered through diatomaceous earth, concentrated *in vacuo* and purified by flash chromatography on silica gel (methylene chloride:methanol:ammonium hydroxide, 96:4:0; then 96:4:1) to give the title compound (676 mg).

20 Step 3.2E 3-[2-[benzyl-(4-pyridin-3-yl-butyl)amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-ylamine

To a stirred solution of benzyl-{2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl}-(4-pyridin-3-yl-butyl)amine (350 mg in 30 mL absolute ethanol) was added ca. 30 mg of Raney[®] nickel. The reaction flask was fitted with a hydrogen balloon, evacuated and recharged with hydrogen (3 times) and stirred at room temperature. After 3 hours the reaction was flushed with nitrogen, filtered over diatomaceous earth and concentrated *in vacuo*. Purification by flash chromatography on silica gel (methylene chloride:methanol:ammonium hydroxide, 96:4:1) gave the title compound (246 mg).

Step 3.2F Benzyl-{2-[2-(3,5-dimethylphenyl)-5-isocyanato-1H-indol-3-yl]ethyl}-(4-pyridin-3-yl-butyl)amine

To a solution of 3-[2-[benzyl-(4-pyridin-3-yl-butyl)amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-ylamine (120 mg in

8 mL dry methylene chloride) at 0 °C was added 26.6 mg triphosgene followed by 0.050 mL pyridine and the mixture stirred at low temperature. After 50 minutes, the mixture was concentrated *in vacuo* to give the crude title compound (120 mg).

5

Step 3.2G 1-[3-[2-[benzyl-(4-pyridin-3-yl-butyl)-amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-1,4-dihydropyridazin-5-one

To a solution of freshly prepared aluminum azide (0.6 mmol in 6 mL dry tetrahydrofuran) was added 120 mg benzyl-[2-[2-(3,5-dimethylphenyl)-5-isocyanato-1H-indol-3-yl]ethyl]-(4-pyridin-3-yl-butyl)amine and the mixture heated to reflux on an oil bath. After 20 hours, the mixture was cooled to room temperature concentrated poured into a mixture of 1M sodium potassium tartarate and ice, stirred vigorously for 40 minutes then partitioned between ethyl acetate and water. The organic portion was washed successively with 1M sodium potassium tartarate, water and brine then dried over sodium sulfate. Purification of the concentrate by flash chromatography on silica gel (methylene chloride:methanol, 88:12) gave the title compound (58 mg).

20

Step 3.2H 1-[3-[2-[benzyl-(4-pyridin-3-yl-butyl)amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-4-methyl-1,4-dihydropyridazin-5-one

To a solution of 1-[3-[2-[benzyl-(4-pyridin-3-yl-butyl)-amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-1,4-dihydropyridazin-5-one (25 mg in 1.5 mL dry *N,N*-dimethylformamide) at 0 °C was added 13 mg potassium carbonate followed by 0.033 mL of a 10% solution of iodomethane in methylene chloride, and the mixture stirred at low temperature. After 2 hours, the reaction was quenched by the addition of saturated aqueous ammonium chloride and the mixture extracted with ethyl acetate. The organic portion was washed successively with water and brine, dried over sodium sulfate and concentrated *in vacuo*. Purification of the concentrate by flash chromatography on silica gel (methylene chloride:methanol, 95:5) gave the title compound (20 mg).

35

Step 3.2I 1-[2-(3,5-dimethylphenyl)-3-[2-(4-pyridin-3-yl-butylamino)ethyl]-1H-indol-5-yl]4-methyl-1,4-dihydrotetrazol-5-one

To a stirred solution of 1-[3-[2-[benzyl-(4-pyridin-3-yl-butyl)amino]ethyl]-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-4-methyl-1,4-dihydrotetrazol-5-one (20 mg in 4 mL methanol) was added 15 mg of 10% palladium hydroxide on carbon catalyst followed by acetic acid (0.020 mL of a 30% solution in water). The reaction flask was fitted with a hydrogen balloon, evacuated and recharged with hydrogen (3 times) and stirred at room temperature. After 30 minutes the reaction was flushed with nitrogen, filtered over diatomaceous earth and concentrated *in vacuo*. Purification by flash chromatography (methylene chloride:methanol:ammonium hydroxide, 90:6.5:1) gave the title compound (16 mg). $m/e = 496$ (M + H)

PREPARATION OF SYNTHETIC INTERMEDIATES


Step A: 4-chloro-N-methoxy-N-methylbutyramide

To a solution of 4-chlorobutyryl chloride (10.0 g in 200 mL of dry methylene chloride) was added 10.4 g of *N,O*-dimethylhydroxylamine hydrochloride. The mixture was stirred under nitrogen and maintained below 25 °C by cooling in an ice bath as necessary while triethylamine (29.1 mL) was added dropwise over about 20 minutes, resulting in precipitation. After 1.5 hours at room temperature, the mixture was concentrated *in vacuo*. The residue was partitioned between 100 mL of diethyl ether and 100 mL of saturated aqueous sodium bicarbonate solution. The organic layer was washed with an additional 100 mL of saturated sodium bicarbonate, and the aqueous fractions were back-extracted with ether. The combined organic phases were dried over sodium sulfate, filtered, and concentrated *in vacuo* to give 10.5 g (90%) of an oil, which had satisfactory purity by ¹H NMR (CDCl₃). Mass spectrum (PB-NH₃/CI): $m/e = 166$ (M + H).

Step B: 3-chloropropyl 3,5-dimethylphenyl ketone

A solution of 10.2 mL (13.9 g; 72 mmol) 5-bromo-*m*-xylene in 200 mL of anhydrous tetrahydrofuran was stirred under nitrogen at -78 °C as 35.8 mL (84 mmol) of 2.5 M *n*-butyllithium in tetrahydrofuran was added dropwise. After 15 minutes at -78 °C, a solution of 10.0 g (60 mmol) of 4-chloro-*N*-methoxy-*N*-methylbutyramide in 30 mL of anhydrous tetrahydrofuran was added dropwise over 25-30 minutes. The resulting solution was maintained at -78 °C for 45 minutes and then warmed briefly to room temperature. The reaction was quenched by addition of 40 mL of 2 N hydrochloric acid and then partitioned between ethyl acetate and water. The organic phase was washed with saturated aqueous sodium bicarbonate solution and then saturated aqueous sodium chloride solution. The organic solution was dried over sodium sulfate, filtered, and concentrated *in vacuo*. Flash chromatography of the residue afforded 8.91 g (70%) of an oil, which had satisfactory purity by ¹H NMR (CDCl₃).

EXAMPLE 3.3



[2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)-1H-indol-3-yl]ethyl]-(4-pyridin-3-yl-butyl)amine

Step 3.3A Benzyl-[2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl]carbamic acid *tert*-butyl ester

To a solution of benzyl-[2-[2-(3,5-dimethylphenyl)-5-nitro-1H-indol-3-yl]ethyl]amine (EXAMPLE 3.2 StepC, 450 mg in 10 mL

tetrahydrofuran and 3 mL water) at 0 °C was added a solution of 491 mg di-*tert*-butyl dicarbonate followed by 236 mg potassium carbonate and the resulting suspension stirred vigorously at 0 °C. After 50 minutes, the reaction was quenched by the addition of excess saturated aqueous ammonium chloride and the mixture extracted with ethyl acetate. The organic portion was dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash chromatography on silica gel (hexane:ethyl acetate, 3:1) to give the title compound (530 mg).

10 Step 3.3B {2-[5-amino-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl}benzylcarbamic acid *tert*-butyl ester
Prepared essentially as described in EXAMPLE 3.2E starting from benzyl-{2-[2-(3,5-dimethylphenyl)-5-nitro-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester (530 mg) to give the title compound
15 (387 mg).

Step 3.3C Benzyl-{2-[2-(3,5-dimethylphenyl)-5-pentanoylamino-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester
To a solution of {2-[5-amino-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl}benzylcarbamic acid *tert*-butyl ester (200 mg in 10 mL dry methylene chloride) at 0 °C was added 0.18 mL triethylamine followed by the dropwise addition of 0.06 mL valeryl chloride and the mixture stirred at low temperature. After 17 minutes, the reaction was quenched by the addition of saturated aqueous sodium bicarbonate and extracted with
20 ethyl acetate. The organic portion was washed successively with saturated sodium bicarbonate and saturated ammonium chloride then dried over sodium sulfate. Purification of the concentrate by flash chromatography on silica gel (hexane:ethyl acetate, 3:2) gave the title compound (230 mg).

30 Step 3.3D Benzyl-{2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester
To a solution of benzyl-{2-[2-(3,5-dimethylphenyl)-5-pentanoylamino-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester (80 mg
35 in 3 mL dry methylene chloride) was added in order 76.6 mg

triphenylphosphine, 21 mg imidazole, 72 mg zinc azide(pyridine complex) and 0.048 mL diethyl azodicarboxylate and the mixture stirred at room temperature. After 15 hours an additional portion of zinc azide•2 pyridine (29 mg) was added. After another 1 hour reaction time, the pot was cooled to room temperature and the reaction mixture applied directly to a silica gel column for purification by flash chromatography (hexane:methylene chloride:ethyl acetate, 3:4:1; then 2:0:1) to give the title compound (57 mg).

10 Step 3.3E Benzyl-(2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl)ethylamine

To a solution of benzyl-(2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl)-ethyl carbamic acid *tert*-butyl ester (57 mg in 3.5 mL methylene chloride) at 0 °C was added 0.12 mL anisole followed by 0.80 mL trifluoroacetic acid and the mixture stirred at 0 °C. After 1.5 hours, the mixture was concentrated *in vacuo* and the residual acid removed by azeotrope with toluene to give the crude title compound in quantitative yield.


20 Step 3.3F Benzyl-(2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl)ethyl-(4-pyridin-3-yl-butyl)amine

Prepared essentially as described in EXAMPLE 3.2D starting from benzyl-(2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl)ethylamine (58 mg) to give the title compound (43 mg).

25 Step 3.3G [2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl]ethyl-(4-pyridin-3-yl-butyl)amine

Prepared essentially as described in EXAMPLE 3.2I starting from benzyl-(2-[5-(2-butylpentazol-1-yl)-2-(3,5-dimethylphenyl)]-1H-indol-3-yl)ethyl-(4-pyridin-3-yl-butyl)amine (43 mg) to give the title compound (34 mg). *m/e* = 522 (M + H).

EXAMPLE 3.4



{2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1H-indol-3-yl]ethyl-(4-pyridin-3-yl-butyl)amine}

5

Step 3.4A {2-[5-cyano-2-(3,5-dimethylphenyl)-1H-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester

Prepared essentially as described in EXAMPLE 3.3A starting from 3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indole-5-carbonitrile (prepared essentially as described in EXAMPLE 3.2 Step A) to give the title compound (300 mg).

10

Step 3.4B {2-[2-(3,5-dimethylphenyl)-5-(N-hydroxycarbamimidoyl)-1H-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester

15

A solution of {2-[5-cyano-2-(3,5-dimethylphenyl)-1H-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester (300 mg in 5 mL ethanol) was added to a suspension of 725 mg potassium carbonate and 273 mg hydroxylamine hydrochloride in 7 mL ethanol and the whole heated to reflux on an oil bath. After 21 hours, the mixture was cooled to room temperature and filtered to remove solids. The filtrate was concentrated *in vacuo* then partitioned between ethyl acetate and water. The organic portion was washed with water, dried over sodium sulfate and the concentrate purified by flash chromatography on silica gel (methylene chloride:methanol, 92:8) to give the title compound (105 mg).

20

25

Step 3.4C {2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1H-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester

To a stirred solution of isovaleric acid (0.025 mL in 4 mL methylene chloride) was added 1-hydroxybenzotriazole (37.8 mg) and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (43.6 mg) and the reagents allowed to mix for 30 minutes. At this time a solution of {2-[2-(3,5-dimethylphenyl)-5-(*N*-hydroxycarbamimidoyl)-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester (81 mg in 3 mL methylene chloride) was added and the reaction stirred at room temperature. After 2 hours, the mixture was concentrated *in vacuo* and purified by flash chromatography on silica gel (methylene chloride:methanol, 96:4) to give the title compound (81 mg).


Step 3.4D 2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1*H*-indol-3-yl]ethylamine

Prepared essentially as described in EXAMPLE 3.3E starting from {2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1*H*-indol-3-yl]ethyl}carbamic acid *tert*-butyl ester (67 mg) to give the title compound (48 mg).

Step 3.4E {2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1*H*-indol-3-yl]ethyl}-(4-pyridin-3-yl-butyl)amine

To a solution of 2-[2-(3,5-dimethylphenyl)-5-(5-isobutyl-[1,2,4]oxadiazol-3-yl)-1*H*-indol-3-yl]ethylamine (22 mg in 1.5 mL chloroform) at 0 °C was added anhydrous magnesium sulfate (38 mg) followed by 4-(3-pyridyl)-butanal (11 mg) and the mixture stirred at low temperature for 15 minutes. At this time sodium borohydride (3.7 mg in 0.50 mL methanol) was added and the mixture stirred at 0° C. After 30 minutes, the reaction was quenched by the addition of water and the mixture extracted with ethyl acetate. The organic portion was washed successively with saturated potassium carbonate and brine then dried over sodium sulfate. Purification of the concentrate by flash chromatography on silica gel (methylene chloride:methanol, 92:8) gave the title compound (26.5 mg). *m/e* = 522 (M + H)

EXAMPLE 3.5



(2-[2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1H-imidazol-2-yl)-ethyl]-1H-indol-3-yl]-ethyl)-(4-pyridin-4-yl-butyl)-amine

5

Step 3.5A 2-[3-(2-*tert*--butoxycarbonylaminoethyl)-2-(3,5-dimethylphenyl)-1H--indol-5-yl]-2-methyl-propionic acid ethyl ester

Prepared essentially as described in EXAMPLE 3.3A

10 starting from 2-[3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H--indol-5-yl]-2-methylpropionic acid ethyl ester (EXAMPLE 2, 1.13 g) to give the title compound (1.28 g).

15 Step 3.5B 2-[3-(2-*tert*--butoxycarbonylamino-ethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid

To a stirred solution of 2-[3-(2-*tert*--butoxycarbonylaminoethyl)-2-(3,5-dimethylphenyl)-1H--indol-5-yl]-2-methyl-propionic acid ethyl ester (1.28 g in 25 mL ethanol) was added 30 mL of 0.5N sodium hydroxide and the mixture heated to 90 °C on an oil bath. After 30 hours the mixture was concentrated *in vacuo*, diluted with water and extracted with diethyl ether (3x). The aqueous layer was then made acidic by the addition of 0.5N hydrochloric acid and extracted with ethyl acetate. The ethyl acetate layer was washed with water and brine, dried over sodium sulfate and concentrated *in vacuo* to give the crude title compound (1.23 g).

25

Step 3.5C (2-[2-(3,5-dimethylphenyl)-5-[1-(methoxymethyl)carbamoyl]-1-methyl-ethyl]-1H-indol-3-yl)ethyl)carbamic acid *tert*--butyl ester

To a suspension of 2-[3-(2-*tert*--butoxycarbonylamino-ethyl)-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-2-methylpropionic acid (1.23g in 15 mL *N,N*-dimethylformamide) at 0° C was added 608 mg of 1-hydroxybenzotriazole (HOBt), 0.48 mL 4-methylmorpholine and 352 mg of *N,O*-dimethylhydroxylamine hydrochloride and the mixture stirred at low temperature. After 15 minutes, 826 mg 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDC) were added and the mixture warmed to room temperature. The reaction was quenched after 3.5 days by concentration *in vacuo*, resuspending in ethyl acetate and washing sequentially with water, 0.3N sodium bisulfate, water, saturated sodium bicarbonate and brine. The organic portion was dried over sodium sulfate and the concentrate purified by flash chromatography on silica gel (hexane:ethyl acetate, 3:1) to give the title compound (905 mg).

15 Step 3.5D [2-[5-(1,1-dimethyl-2-oxo-ethyl)-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl]carbamic acid *tert*--butyl ester

To a solution of (2-[2-(3,5-dimethylphenyl)-5-[1-(methoxymethylcarbonyl)-1-methyl-ethyl]-1*H*-indol-3-yl]ethyl)carbamic acid *tert*--butyl ester (296 mg in 5 mL dry tetrahydrofuran) at 0 °C was added 1.8 mL of a 1M lithium aluminum hydride solution in tetrahydrofuran and the mixture stirred at low temperature. After 1 hour, the reaction was quenched by the careful addition of 0.3M aqueous sodium bisulfate solution. The resulting mixture was extracted with ethyl acetate and the organic portion washed successively with 0.3M aqueous sodium bisulfate, water and brine. This was then dried over sodium sulfate, concentrated *in vacuo* and purified by flash chromatography on silica gel (hexane:ethyl acetate, 85:15) to give the title compound (253 mg).

30 Step 3.5E (2-[2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1*H*-imidazol-2-yl)ethyl]-1*H*-indol-3-yl]ethyl)carbamic acid *tert*--butyl ester

To a solution of {2-[5-(1,1-dimethyl-2-oxo-ethyl)-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]ethyl}carbamic acid *tert*--butyl ester (475 mg in 15 mL methanol) was added 1 mL of a 40% aqueous solution of

pyruvic aldehyde followed by 2.2 mL ammonium hydroxide and the mixture stirred at room temperature. After 2 days, an additional portion of 40% aqueous pyruvic aldehyde (0.50 mL) and ammonium hydroxide (1.1 mL) were added. Finally after 4 days the reaction mixture was concentrated *in vacuo*, the residue resuspended in ethyl acetate and washed sequentially with water and brine. The combined organics were dried over sodium sulfate and the concentrate purified by flash chromatography on silica gel (hexane:ethyl acetate, 1:4) to give the title compound (402 mg).

10

Step 3.5F 2-{2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1H-imidazol-2-yl)ethyl]-1H-indol-3-yl}ethylamine

Prepared essentially as described in EXAMPLE 3.3E starting from (2-{2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1H-imidazol-2-yl)ethyl]-1H-indol-3-yl}ethyl)carbamic acid *tert*-butyl ester (353 mg) to give the title compound (198 mg).

15


Step 3.5G (2-{2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1H-imidazol-2-yl)-ethyl]-1H-indol-3-yl}-ethyl)-(4-pyridin-4-yl-butyl)-amine

20

Prepared essentially as described in EXAMPLE 3.2D starting from 2-{2-(3,5-dimethylphenyl)-5-[1-methyl-1-(4-methyl-1H-imidazol-2-yl)ethyl]-1H-indol-3-yl}ethylamine (97 mg) and using 4-pyridin-4-yl butyraldehyde to give the title compound (73 mg). *m/e* = 520 (M+1)

25

EXAMPLE 4



5 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-(1,1-dimethyl-4-pyridin-4-yl-butylamino)-ethyl]-1H-indol-5-yl]-2-methylpropan-1-one

Step 4A 1-(3-methyl-but-2-enyl)-tetrahydrothiophenium bromide

To a solution of prenyl bromide (2.9 g in 10 mL dry tetrahydrofuran) at 0 °C was added 1.8 mL tetrahydrothiophene and the mixture allowed to warm to room temperature. After 22 hours, the mixture was concentrated *in vacuo* and the remaining starting materials removed by azeotrope with toluene to give the crude title compound as a white solid (2.2 g).

15 Step 4B 4-[3-(2-methylpropenyl)oxiranyl]pyridine

To a suspension of 1-(3-methyl-but-2-enyl)-tetrahydrothiophenium bromide (381 mg in 5 mL dry tetrahydrofuran) at 0 °C was added 0.31 mL pyridine-4-carboxaldehyde followed by 111 mg sodium hydride and the mixture allowed to warm to room temperature. After 1.5 hours, an additional 0.15 mL pyridine-4-carboxaldehyde was added and after 30 minutes the reaction quenched by the addition of water. The mixture was partitioned between ethyl acetate and water and the organic portion washed with saturated aqueous sodium bicarbonate and dried over sodium sulfate. Purification of the concentrate by flash chromatography on silica gel (hexane:ethyl acetate, 1:2) gave the title compound (138 mg).


Step 4C 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-{2-(3,5-dimethyl-phenyl)-3-[2-(4-hydroxy-1,1-dimethyl-4-pyridin-4-yl-but-2-enylamino)ethyl]-1H-indol-5-yl}-2-methylpropan-1-one

To a solution of 2-[3-(2-aminoethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-1-(7-azabicyclo[2.2.1]hept-7-yl)-2-methylpropan-1-one (prepared essentially as described in EXAMPLE 2, 70 mg in 6 mL dry tetrahydrofuran) was added 2 mL of a solution of 86 mg 4-[3-(2-methylpropenyl)oxiranyl]pyridine in tetrahydrofuran followed by 24 mg tetrakis(triphenylphosphine)palladium and the mixture heated to 65 °C on an oil bath. After 2 hours, the mixture was cooled to room temperature, concentrated in vacuo. The residue was purified by flash chromatography on silica gel (methylene chloride:methanol, 92:8; then 88:12) to give the title compound (91 mg).

Step 4D 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-{2-(3,5-dimethyl-phenyl)-3-[2-(1,1-dimethyl-4-pyridin-4-yl-butylamino)-ethyl]-1H-indol-5-yl}-2-methyl-propan-1-one

To a solution of 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-{2-(3,5-dimethylphenyl)-3-[2-(4-hydroxy-1,1-dimethyl-4-pyridin-4-yl-but-2-enylamino)ethyl]-1H-indol-5-yl}-2-methylpropan-1-one (27 mg in a mixture of 2 mL dry tetrahydrofuran and 2 mL ethyl acetate) at 0 °C was added 0.010 mL trifluoroacetic anhydride followed by 0.009 mL triethylamine and the mixture stirred at low temperature. After 15 minutes, 0.030 mL acetic acid was added and along with 28 mg palladium hydroxide on carbon. The reaction flask was fitted with a hydrogen balloon, evacuated and recharged with hydrogen (3 times) and stirred at room temperature. After 8 hours the reaction was flushed with nitrogen, filtered over diatomaceous earth and concentrated *in vacuo*. Purification by flash chromatography on silica gel (methylene chloride:methanol:ammonium hydroxide, 94:6:1) gave the title compound (15 mg). *m/e* = 591 (M+H).

EXAMPLE 5.1



1-(7-azabicyclo[2.2.1]hept-7-yl)-2-{2-(3,5-dimethylphenyl)-3-[1-methyl-2-(4-pyridin-4-yl-butylamino)ethyl]-1H-indol-5-yl}-2-methylpropan-1-one

5

Step 5.1A 2-methylcyclopropanecarboxylic acid N-methoxy-N-methyl-amide

To a solution of 2-methylcyclopropanecarboxylic acid (10 g in a mixture of 200 mL benzene and 2 mL *N,N*-dimethylformamide) at 0° C was added 10.5 mL of oxalyl chloride and the mixture stirred at 0° C for 30 minutes then warmed to room temperature for 30 minutes. At this time, 14.6 g of *N,O*-dimethylhydroxylamine hydrochloride was added followed by 41 mL of triethylamine. The mixture was stirred at room temperature for one hour then quenched by the addition of saturated sodium bicarbonate. The aqueous portion was extracted with ethyl acetate and the combined organics washed with brine, dried over sodium sulfate and concentrated *in vacuo*. The product was purified by distillation under reduced pressure to give 8.9 g as an oil.

20 Step 5.1B (3,5-dimethylphenyl)-(2-methylcyclopropyl)methanone

To a solution of 5-bromo-*meta*-xylene (5.7 mL in 120 mL of dry tetrahydrofuran) at -78° C was added 30.6 mL of a 1.4M solution of *n*-butyllithium in hexane and the mixture stirred at low temperature. After 15 minutes, a solution of 2-methylcyclopropanecarboxylic acid *N*-methoxy-*N*-methyl-amide (5.0 g in 50 mL tetrahydrofuran) was added dropwise over 5 minutes and the mixture then allowed to warm slowly to room temperature. After 1 hour, the reaction was quenched by the addition of 20 mL 2N hydrochloric acid and 40 mL water. This was extracted with ethyl acetate washed with saturated sodium bicarbonate

and brine then dried over sodium sulfate to give 6.95 g of the title compound (crude).

5 Step 5.1C 2-[3-(2-amino-1-methylethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid ethyl ester

To a solution of 2-(4-hydrazinophenyl)-2-methylpropionic acid ethyl ester (5.7 g in 20 mL n-butanol) was added 4 g (3,5-dimethylphenyl)-(2-methylcyclopropyl)methanone followed by 1.3 mL conc. hydrochloric acid and the mixture heated to 110° C on an oil bath. After 16 hours, the mixture was cooled to room temperature and concentrated *in vacuo*. The residue was dissolved in ethyl acetate and washed sequentially with 0.5 N sodium hydroxide, water and brine. The combined organics were dried over sodium sulfate and concentrated *in vacuo*. Purification by flash chromatography on silica gel (methylene chloride:methanol, 95:5) gave the title compound (2.5 g).

20 Step 5.1D 2-{2-(3,5-dimethylphenyl)-3-[1-methyl-2-(4-pyridin-4-yl-butylamino)-ethyl]-1H-indol-5-yl}-2-methylpropionic acid ethyl ester

Prepared essentially as described in EXAMPLE 2, Step A from 2-[3-(2-amino-1-methylethyl)-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid ethyl ester (233 mg) to give the title compound (258 mg).

25 Step 5.1E 2-[3-{2-[benzyloxycarbonyl-(4-pyridin-4-yl-butyl)-amino]-1-methyl-ethyl}-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid ethyl ester

Prepared essentially as described in EXAMPLE 2, Step B from 2-{2-(3,5-dimethylphenyl)-3-[1-methyl-2-(4-pyridin-4-yl-butylamino)-ethyl]-1H-indol-5-yl}-2-methylpropionic acid ethyl ester (258 mg) to give the title compound (240 mg).

35 Step 5.1F 2-[3-{2-[benzyloxycarbonyl-(4-pyridin-4-yl-butyl)amino]-1-methylethyl}-2-(3,5-dimethylphenyl)-1H-indol-5-yl]-2-methylpropionic acid

Prepared essentially as described in EXAMPLE 2, Step C from 2-[3-{2-[benzyloxycarbonyl-(4-pyridin-4-yl-butyl)-amino]-1-methylethyl}-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-2-methylpropionic acid ethyl ester (240 mg) to give the title compound (222 mg).

5

Step 5.1G {2-[5-[2-(7-azabicyclo[2.2.1]hept-7-yl)-1,1-dimethyl-2-oxo-ethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]propyl}-(4-pyridin-4-yl-butyl)carbamic acid benzyl ester

Prepared essentially as described in EXAMPLE 2, Step D from 2-[3-{2-[benzyloxycarbonyl-(4-pyridin-4-yl-butyl)amino]-1-methylethyl}-2-(3,5-dimethylphenyl)-1*H*-indol-5-yl]-2-methylpropionic acid (91 mg) to give the title compound (66 mg).

10


Step 5.1H 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-{2-(3,5-dimethylphenyl)-3-[1-methyl-2-(4-pyridin-4-yl-butylamino)ethyl]-1*H*-indol-5-yl}-2-methylpropan-1-one

15

Prepared essentially as described in EXAMPLE 2, Step E from {2-[5-[2-(7-azabicyclo[2.2.1]hept-7-yl)-1,1-dimethyl-2-oxo-ethyl]-2-(3,5-dimethylphenyl)-1*H*-indol-3-yl]propyl}-(4-pyridin-4-yl-butyl)carbamic acid benzyl ester (62 mg) to give the title compound (27 mg). *m/e* = 577 (*M* + *H*).

20

EXAMPLE 5.2




25

1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[methyl-[4-(pyridin-4-yl)butylamino]ethyl]-1H-indol-5-yl]-2-methylpropan-1-one













5 A dry flask containing 120 mg (0.21 mmol) of 1-(7-azabicyclo[2.2.1]hept-7-yl)-2-[2-(3,5-dimethylphenyl)-3-[2-[4-(pyridin-4-yl)butylamino]ethyl]-1H-indol-5-yl)-2-methylpropan-1-one (prepared essentially as described in EXAMPLE 2), 63.0 mg (2.1 mmol) of paraformaldehyde, and 200 mg of powdered 3A molecular sieves was
10 fitted with a septum and purged thoroughly with nitrogen. Next, 5 mL of methanol and 0.121 mL (126.1 mg, 2.1 mmol) of glacial acetic acid were added, and the mixture was stirred at room temperature for 15 minutes. Then 52.8 mg (0.84 mmol) of sodium cyanoborohydride was added, followed after an additional 25 minutes by 2.5 mL of anhydrous
15 tetrahydrofuran. After 1 day, the mixture was filtered, and the filter cake was washed thoroughly with methylene chloride. The filtrate was shaken in a separatory funnel with water. The aqueous phase was extracted an additional 3 times with methylene chloride. The combined organic fractions were dried over sodium sulfate, filtered, and
20 concentrated *in vacuo*. The residue was flash chromatographed on silica gel (gradient elution with 99:1:0.1 to 95:5:0.5 CH₂Cl₂-MeOH-concd. NH₄OH) to yield 95.4 mg (79%) of a yellow, stiff foam; homogeneous by TLC in 95:5:0.5 CH₂Cl₂-MeOH-concd. NH₄OH. 500 MHz ¹H NMR (CDCl₃) was consistent with the assigned structure. Mass spectrum
25 (ESI): *m/e* = 577.5 (M + H).












EXAMPLE 6













30 Following procedures similar to that described in EXAMPLES 1 through 5, and Schemes O and P, the following compounds are prepared:



























Ex. #	-(A)-R ₁	R ₂
6A		H
6B		Me
6C		H
6D		H
6E		Me
6F		H
6G		H
6H		Me












6I		H
6J		H
6K		Me
6L		H
6M		Me
6N		H
6O		H
6P		H
6Q		H
6R		H
6S		Me
6T		H












6U		H
6V		H
6W		H
6X		H
6Y		H
6Z		H
6AA		H
6BB		H
6CC		Me
6DD		H
6EE		Me












6FF		H
6GG		H
6HH		H
6II		Me
6JJ		i-Pr
6KK		Et
6LL		H
6MM		H
6NN		H
6OO		H
6PP		H
6QQ		H











6RR		H
6SS		H
6TT		H
6UU		H
6VV		H
6WW		H
6XX		H
6YY		H
6ZZ		H
6AAA		
6BBB		H
6CCC		H

6DDD		H
6EEE		H
6FFF		H
6GGG		H
6HHH		H
6III		H
6JJJ		H
6KKK		H
6LLL		H
6MMM		H
6NNN		H
6OOO		H

6PPP		H
6QQQ		H
6RRR		H
6SSS		H
6TTT		H
6UUU		H
6VVV		H
6WWW		H
6XXX		H
6YYY		H
6ZZZ		H


6AAAA		H
6BBBB		H
6CCCC		H
6DDDD		H
6EEEE		H
6FFFF		H
6GGGG		H
6HHHH		H
6IIII		H
6JJJJ		H
6KKKK		H

6LLLL		H
6MMMM		H
6NNNN		H
60000		H
6PPPP		H
6QQQQ		H
6RRRR		H
6SSSS		H
6TTTT		H
6UUUU		H
6VVVV		H







6WWWW		H
6XXXX		H
6YYYY		H
6ZZZZ		H
6AAAA		H
6BBBB		H
6CCCC		H
6DDDD		H
6EEEE		H
6FFFF		H

EXAMPLE 7

Following procedures similar to that described in
 EXAMPLES 1 through 5, and Scheme P, the following compounds are
 5 prepared:




Ex. #	X-R7R8	-(A)-R1	R2	R9, R9a
7A			H	H,H
7B			H	H,H
7C			H	H,H
7D			H	H,H
7E			H	Me, Me



7F			H	H,H
7G			H	H,H
7H			H	H,H












EXAMPLE 8








Following procedures similar to that described in
5 EXAMPLES 1 through 5, and Scheme P, the following compounds are
prepared:



10


Ex. #	-(A)-R ₁	R ₂
8A		H
8B		Me

8C		Et
8D		i-Bu
8E		i-Pr
8F		H
8G		H
8H		H
8I		H
8J		H
8K		H
8L		H
8M		H









8N		H
8O		H
8P		H
8Q		H
8R		H
8S		H
8T		H

EXAMPLE 9

- 5 Following procedures similar to that described in
EXAMPLES 1 through 5, and Scheme P, the following compounds are
prepared:




Ex. #	X-R7R8	n
9A		4
9B		4
9C		4
9D		4
9E		4
9F		4

9G		4
9H		2
9I		4
9J		4
9K		4
9L		4
9M		4
9N		2

EXAMPLE 10

Following procedures similar to that described in
EXAMPLES 1 through 5, and Scheme P, the following compounds are
prepared:



5


Ex. #	R
10A	

EXAMPLE 11












10












Following procedures similar to that described in
EXAMPLES 1 through 5, and Scheme P, the following compounds are
prepared:












15
















Ex. #	-(A)-R ₁	R ₂
-------	---------------------	----------------

11A		H
11B		H
11C		H
11D		H
11E		H
11F		H
11G		H
11H		H
11I		H
11J		H
11K		H

11L		H
11M		H
11N		H
11O		H
11P		H
11Q		H
11R		H
11S		H
11T		H
11U		H
11V		H

11W		H
11X		H
11Y		H
11Z		H
11AA		H
11BB		H
11CC		H
11DD		H
11EE		H
11FF		H
11GG		H

11HH	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11II	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11JJ	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11KK	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11LL	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11MM	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11NN	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11OO	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11PP	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11QQ	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11RR	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H
11SS	 <chem>CC1=CC(=C(C=C1)C)C(=CC1=CC=C1O1=CC=CC=C1)OCC</chem>	H


11TT		H
11UU		H
11VV		H
11WW		H


EXAMPLE 12










5

Following procedures similar to that described in EXAMPLES 1 through 5, and Scheme P, the following compounds are prepared:

10




Ex. #	-(A)-R ₁	R ₂
12A		H















12B		H
12C		H
12D		H
12E		H
12F		
12G		H
12H		
12I		H
12J		H

EXAMPLE 13

5 Following procedures similar to that described in
EXAMPLES 1 through 5, and Scheme P, the following compounds are
prepared:




Ex. #	-(A)-R ₁	-NR
13A		
13B		
13C		
13D		
13E		
13F		
13G		
13H		
13I		

13J		
13K		
13L		
13M		
13N		
13O		
13P		

WHAT IS CLAIMED IS:

1. A compound of the formula


5



wherein

A is C1-C6 alkyl, substituted C1-C6 alkyl, C3-C7 cycloalkyl, substituted C3-C7 cycloalkyl, C3-C6 alkenyl, substituted C3-C6 alkenyl, C3-C6 alkynyl, substituted C3-C6 alkynyl, C1-C6 alkoxy, or C0-C5 alkyl-S(O)_n-C0-C5 alkyl, C0-C5 alkyl-O-C0-C5 alkyl, C0-C5 alkyl-NR₁₈-C0-C5 alkyl where R₁₈ and the

10




C0-C5 alkyl can be joined to form a ring, or a single bond;

15

R₀ is hydrogen, C1-C6 alkyl, substituted C1-C6 alkyl, wherein the substituents are as defined below; aryl, substituted aryl, aralkyl or substituted aralkyl, wherein the substituents are as defined for R₃, R₄ and R₅;

20 R₁ is



the nitrogen atoms contained in the R₁ heteroaromatic rings may exist

either as drawn or, when chemically allowed, in their oxidized (N→O, N-OH) state;

- 5 R₂ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aralkyl, substituted aralkyl, aryl, substituted aryl, alkyl -OR₁₁, C₁-C₆(NR₁₁R₁₂), C₁-C₆(CONR₁₁R₁₂) or C(NR₁₁R₁₂)NH;

R₂ and A taken together form a ring of 5-7 atoms;

- 10 R₃, R₄ and R₅ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, CN, nitro, C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, aryl,

- substituted aryl, aralkyl, substituted aralkyl, $R_{11}O(CH_2)_p$ -, $R_{11}C(O)O(CH_2)_p$ -, $R_{11}OC(O)(CH_2)_p$ -, $-(CH_2)_pS(O)_nR_{17}$ -, $-(CH_2)_pC(O)NR_{11}R_{12}$ or halogen; wherein R_{17} is hydrogen, C₁-C₆ alkyl, C₁-C₃ perfluoroalkyl, aryl or substituted aryl;
- 5 R_3 and R_4 taken together form a carbocyclic ring of 3-7 carbon atoms or a heterocyclic ring containing 1-3 heteroatoms selected from N, O and S;
- R_6 is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, C₁-C₃ perfluoroalkyl, CN, NO₂, halogen,
- 10 $R_{11}O(CH_2)_p$ -, $NR_{21}C(O)R_{20}$, $NR_{21}C(O)NR_{20}R_{21}$ or SO_nR_{20} ;
- R_7 is hydrogen, C₁-C₆ alkyl, or substituted C₁-C₆ alkyl, unless X is hydrogen or halogen, then R_7 is absent;
- R_8 is $C(O)OR_{20}$, $C(O)NR_{20}R_{21}$, $NR_{20}R_{21}$, $C(O)R_{20}$, $NR_{21}C(O)R_{20}$, $NR_{21}C(O)NR_{20}R_{21}$, $NR_{20}S(O)_2R_{21}$, $NR_{21}S(O)_2NR_{20}R_{21}$,
- 15 $OC(O)R_{20}$, $OC(O)NR_{20}R_{21}$, OR_{20} , SO_nR_{20} , $S(O)_nNR_{20}R_{21}$, a heterocyclic ring or bicyclic heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R_3 , R_4 and R_5 , C₁-C₆ alkyl or substituted C₁-C₆ alkyl; or
- 20 R_7 and R_8 taken together form a heterocyclic ring containing one or more heteroatoms selected from N, O or S which can be optionally substituted by R_3 , R_4 and R_5 ;
- R_9 and R_{9a} are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl; aryl or substituted aryl, aralkyl or substituted aralkyl
- 25 when $m \neq 0$; or
- R_9 and R_{9a} taken together form a carbocyclic ring of 3-7 atoms or $\overset{O}{\parallel}$ when $m \neq 0$;
- R_9 and A taken together form a heterocyclic ring containing 3-7 carbon atoms and one or more heteroatoms when $m \neq 0$; or
- 30 R_{10} and R_{10a} are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, aralkyl or substituted aralkyl; or


- R₁₀ and R_{10a} taken together form a carbocyclic ring of 3-7 atoms or $\text{O}=\text{C}$;
 R₉ and R₁₀ taken together form a carbocyclic ring of 3-7 carbon atoms or
 a heterocyclic ring containing one or more heteroatoms
 when m≠0; or
- 5 R₉ and R₂ taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms when m≠0; or
 R₁₀ and R₂ taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms;
- 10 R₁₀ and A taken together form a heterocyclic ring containing 3-7 carbon
 atoms and one or more heteroatoms; or
 R₁₁ and R₁₂ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-
 C₆ alkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, a
 carbocyclic ring of 3-7 atoms or a substituted carbocyclic ring
 containing 3-7 atoms;
- 15 R₁₁ and R₁₂ taken together can form an optionally substituted ring of 3-7
 atoms;
- R₁₃ is hydrogen, OH, NR₇R₈, NR₁₁SO₂(C₁-C₆ alkyl),
 NR₁₁SO₂(substituted C₁-C₆ alkyl), NR₁₁SO₂(aryl),
 NR₁₁SO₂(substituted aryl), NR₁₁SO₂(C₁-C₃ perfluoroalkyl);
 20 SO₂NR₁₁(C₁-C₆ alkyl), SO₂NR₁₁(substituted C₁-C₆ alkyl),
 SO₂NR₁₁(aryl), SO₂NR₁₁(substituted aryl), SO₂NR₁₁(C₁-C₃
 perfluoroalkyl); SO₂NR₁₁(C(O)C₁-C₆ alkyl); SO₂NR₁₁(C(O)-
 substituted C₁-C₆ alkyl); SO₂NR₁₁(C(O)-aryl);
 SO₂NR₁₁(C(O)-substituted aryl); S(O)_n(C₁-C₆ alkyl); S(O)_n
 25 (substituted C₁-C₆ alkyl), S(O)_n(aryl), S(O)_n(substituted
 aryl), C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, C₁-C₆
 alkoxy, substituted C₁-C₆ alkoxy, COOH, halogen, NO₂ or
 CN;
- 30 R₁₄ and R₁₅ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆
 alkyl, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, CN, nitro,
 C₁-C₃ perfluoroalkyl, C₁-C₃ perfluoroalkoxy, aryl,
 substituted aryl, aralkyl, substituted aralkyl, R₁₁O(CH₂)_p-,
 R₁₁C(O)O(CH₂)_p-, R₁₁OC(O)(CH₂)_p-, -(CH₂)_pS(O)_nR₁₇, -

- (CH₂)_pC(O)NR₁₁R₁₂ or halogen; wherein R₁₇ is hydrogen, C₁-C₆ alkyl, C₁-C₃ perfluoroalkyl, aryl or substituted aryl;
- R₁₆ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, or N(R₁₁R₁₂);
- 5 R₁₈ is hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, C(O)OR₁₁, C(O)NR₁₁R₁₂, C(O)R₁₁, S(O)_nR₁₁;
- R₁₉ is either the definition of R₁₃ or R₁₄;
- R₂₀ and R₂₁ are independently hydrogen, C₁-C₆ alkyl, substituted C₁-C₆ alkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, a
- 10 carbocyclic ring of 3-7 atoms, a substituted carbocyclic ring containing 3-7 atoms, a heterocyclic ring or bicyclic heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅, C₁-C₆-alkyl substituted by a heterocyclic ring or bicyclic
- 15 heterocyclic ring with from 1 to 4 heteroatoms selected from N, O or S which can be optionally substituted by R₃, R₄ and R₅;
- R₂₀ and R₂₁ taken together can form an optionally substituted ring of 3-7 atoms;
- 20 X is N, O, S(O)_n, C(O), (CR₁₁R₁₂)_p, a single bond to R₈, C₂-C₆ alkenyl, substituted C₂-C₆ alkenyl, C₂-C₆ alkynyl, or substituted C₂-C₆ alkynyl; when X is O, S(O)_n, C(O), or CR₁₁R₁₂ only R₈ is possible;
- Z is O, S or NR₁₁;
- 25 m is 0-3;
- n is 0-2;
- p is 0-4; and
- 30 the alkyl, cycloalkyl, alkenyl and alkynyl substituents are selected from C₁-C₆ alkyl, C₃-C₇ cycloalkyl, aryl, substituted aryl, aralkyl, substituted aralkyl, hydroxy, oxo, cyano, C₁-C₆ alkoxy, fluoro, C(O)OR₁₁, aryl C₁-C₃ alkoxy, substituted aryl C₁-C₃ alkoxy, and the aryl substituents are as defined for R₃, R₄ and R₅;

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof.

5

2. The compound according to Claim 1 of the formula

























10 or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;













wherein R_1 , R_2 and A are as indicated in the table below:













15














$-(A)-R_1$	R_2
	H
	Me
	H
	H











	Me
	H
	H
	Me
	H
	H
	Me
	H
	Me
	H
	H













	H
	H
	H
	Me
	H
	H
	H
	H
	H
	H
	H












	H
	H
	Me
	H
	Me
	H
	H
	H
	Me
	i-Pr
	Et
	H












	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H




	H
	H
	
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H


	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H



3. The compound according to Claim 1 of the formula

















5


or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;

10 wherein R_1 , R_2 , R_9 , R_{9a} , $X-R_7R_8$ and A are as indicated in the table below:

$X-R_7R_8$	$-(A)-R_1$	R_2	R_9, R_{9a}
		H	H,H

		H	H,H
		H	H,H
		H	H,H
		H	Me, Me
		H	H,H
		H	H,H
		H	H,H

4. The compound according to Claim 1 of the formula
















or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;

5


wherein R_1 , R_2 and A are as indicated in the table below:

$-(A)-R_1$	R_2
	H
	Me
	Et
	i-Bu
	i-Pr
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H


5. The compound according to Claim 1 of the formula











5

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;


10 wherein X-R7R8 and n are as indicated in the table below:

X-R7R8	n
	4

	4
	4
	4
	4
	4
	4
	2
	4


	4
	4
	4
	4
	2

6. The compound according to Claim 1 of the formula




5

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;



wherein R is as indicated in the table below:












- 5 7. The compound according to Claim 1 of the formula
























or a pharmaceutically acceptable addition salt and/or hydrate thereof, or
 where applicable, a geometric or optical isomer or racemic mixture
 10 thereof;













wherein R₁, R₂ and A are as indicated in the table below:



-(A)-R ₁	R ₂
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H


	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H
	H

	H
	H

8. The compound according to Claim 1 of the formula













5

or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;


10

wherein R_1 , R_2 and A are as indicated in the table below:

$-(A)-R_1$	R_2
	H
	H
	H





















	H
	H
	
	H
	
	H
	H













9. The compound according to Claim 1 of the formula



or a pharmaceutically acceptable addition salt and/or hydrate thereof, or where applicable, a geometric or optical isomer or racemic mixture thereof;

5 wherein R_1 , NR and A are as indicated in the table below:

-(A)- R_1	-NR
	
	
	
	
	
	
	
	
	
	

10. A pharmaceutical composition which comprises an effective amount of a compound as defined in Claim 1 and a pharmaceutically acceptable carrier therefor.

5

11. A method for antagonizing gonadotropin-releasing hormone in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1 to a subject suffering from a gonadotropin-releasing hormone derived disorder.

10

12. A method according to Claim 11 wherein the gonadotropin-releasing hormone derived disorder is a sex-hormone related condition.

5 13. A method according to Claim 11 wherein the gonadotropin-releasing hormone derived disorder is a sex hormone dependent cancer, benign prostatic hypertrophy or myoma of the uterus.

10 14. A method according to Claim 13 wherein the sex hormone dependent cancer is selected from the group consisting of prostatic cancer, uterine cancer, breast cancer and pituitary gonadotrophe adenomas.

15 15. A method according to Claim 12 wherein the sex hormone related condition is selected from the group consisting of endometriosis, polycystic ovarian disease, uterine fibroids and precocious puberty.

20 16. A method for preventing pregnancy in a subject in need thereof which comprises administering an effective amount of a compound as defined in Claim 1.

25 17. A method for treating lupus erythematosus in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1.

30 18. A method for treating irritable bowel syndrome in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1.

19. A method for treating premenstrual syndrome in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1.

20. A method for treating hirsutism in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1.

5 21. A method for treating short stature or a growth hormone deficiency in a subject in need thereof which comprises administering to said subject an effective amount of a compound which stimulates the endogenous production or release of growth hormone and an effective amount of a compound as defined in Claim 1.

10 22. A method for treating sleep disorders in a subject in need thereof which comprises administering to said subject an effective amount of a compound as defined in Claim 1.

15 23. The method of Claim 22 wherein the sleep disorder is sleep apnea.

20 24. A pharmaceutical composition which comprises an inert carrier and an effective amount of a compound which stimulates the endogenous production or release of growth hormone in combination with a compound as defined in Claim 1.

25 25. A pharmaceutical composition made by combining the compound of Claim 1 and a pharmaceutically acceptable carrier therefor.

30 26. A process for making a pharmaceutical composition comprising combining a compound of Claim 1 and a pharmaceutically acceptable carrier.

27. A pharmaceutical comprising a compound having luteinizing hormone releasing hormone activity in combination with a compound of Claim 1.

28. The pharmaceutical of Claim 27 wherein the compound having luteinizing hormone releasing hormone activity is a peptide compound.

5 29. The pharmaceutical of Claim 28 wherein the peptide compound is a natural hormone or an analog thereof.

 30. The pharmaceutical of Claim 28 wherein the peptide
compound is a compound selected from the group consisting of
10 leuprorelin, gonadorelin, buserelin, triptorelin, goserelin, nafarelin,
histrelin, deslorelin, meterelin and recirelin.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/15581

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/14 A61K31/40 C07D401/12 C07D413/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	S.S.TIWARI ET AL.: "New indolic anti-tremor agents" PHARMAZIE., vol. 35, no. 12, - 1980 pages 742-743, XP002121006 VEB VERLAG VOLK UND GESUNDHEIT. BERLIN., DD ISSN: 0031-7144 * Table 3: compounds 11-22 *	1
X	WO 97 21703 A (MERCK & CO., INC.) 19 June 1997 (1997-06-19) cited in the application * complete document *	1,10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

4 November 1999

Date of mailing of the international search report

16/11/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Van Bijlen, H

INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 99/15581

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 97 21704 A (MERCK & CO., INC.) 19 June 1997 (1997-06-19) cited in the application * complete document *	1,10
P,X	WO 99 21533 A (MERCK & CO., INC.) 6 May 1999 (1999-05-06) * complete document *	1,10

INTERNATIONAL SEARCH REPORT

national application No.

PCT/US 99/ 15581

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11-19
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 11-19
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 99/15581

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9721703 A	19-06-1997	AU 709090 B	19-08-1999
		AU 1281997 A	03-07-1997
		CA 2240048 A	19-06-1997
		CN 1208413 A	17-02-1999
		CZ 9801832 A	16-09-1998
		EP 0871624 A	21-10-1998
		JP 11501329 T	02-02-1999
		NO 982731 A	13-08-1998
		PL 327132 A	23-11-1998
		SK 77598 A	11-01-1999
		US 5849764 A	15-12-1998

WO 9721704 04 A		NONE	

WO 9921533 A	06-05-1999	AU 1198599 A	17-05-1999
